



**MEMORANDUM OF UNDERSTANDING  
ON THE CONSERVATION AND  
MANAGEMENT OF MARINE TURTLES  
AND THEIR HABITATS OF THE INDIAN  
OCEAN AND SOUTH-EAST ASIA**

CMS/IOSEA/MOS8/Inf.11.3

19 October 2019

Original: English

8<sup>TH</sup> MEETING OF THE SIGNATORY STATES

Da Nang, Viet Nam, 21-25 October 2019

Agenda Item 11.3

**GUIDANCE ON HATCHERIES**

*(Prepared by the Secretariat)*

Action requested:

- Take note of the guidance developed externally for the Northern Indian Ocean Sub-Region

## GUIDANCE ON HATCHERIES

1. Hatcheries for marine turtles are common in several countries in the IOSEA region, including in Signatory States of the IOSEA Marine Turtle MOU.
2. Most recently, the approximate number of hatcheries and/or their practices have been described for the following Signatory States:

### *Northern Indian Ocean Sub-region*

Bangladesh: ≤ 33. (Phillott. 2018. IOTN 27: 29-30.)

India: >31 (Phillott et al. 2019. In Press.)

Sri Lanka: ≥11 (Phillott et al. 2018. IOTN 27: 3-17.)

Pakistan: 3 (Phillott et al. 2018. IOTN 27: 2-8.)

### *North-western Indian Ocean Sub-region*

Iran- ≥1 (Pazira et al. 2016. Reg.Stud.Mar.Sci. 3: 216-224.)

Qatar- ≥1. (Chatting et al. 2008, PLoS ONE 13: e0203257.)

### *South-East Asia + Sub-region*

Indonesia- ≥3 (Maulany et al. 2012. Copeia 2012: 738-747; Tapilatu & Tiwari. 2007. Chel.Conserv.Biol. 6: 154-158; Tapilatu et al. 2017. Biodiversitas 18: 129-136.)

Myanmar- ≥1. (Lwin. 2009. IOTN 10: 14-18.)

Hatcheries have been previously described in the following Signatory States, but the current number and practices are unknown: Australia, Kenya, Malaysia, Maldives (for head-starting of hatchlings only), Papua New Guinea, Philippines, Qatar, Singapore, Thailand, United Republic of Tanzania and Viet Nam.

3. Hatcheries can be a very valuable conservation tool, if natural nesting conditions are threatened by issues such as flooding, erosion, predation or heavy poaching, and these cannot be brought under control by other means. However, successful hatching of turtles requires very specific handling and conditions, and these are not easy to meet, which may have adverse impacts on success rates. Further, there is a danger of misuse of such facilities to take advantage of paying tourists and volunteers, rather than benefit turtles.
4. At the 7<sup>th</sup> Meeting of Signatory States, the need for assess hatchery management practices was discussed in connection with the Provisional List of Project Concepts Resulting from the Updated (2012) Leatherback Assessment. In addition, capacity-building needs regarding hatchery management were identified in the NIO sub-region, and advice on hatchery management practice requested for e.g. the Maldives and Sri Lanka. In addition, it was recommended that Hawksbill hatchery production success be assessed in Indonesia and the Philippines.
5. In line with these recommendations, the Northern Indian Ocean Marine Turtle Task Force (NIO-MTTF) included in its work plan agreed at its second meeting (2018) plans to “[d]evelop/identify a standardized monitoring protocol to guide and collect data on hatchery and head-starting practices”.

6. The attached paper on *Best Practices for Sea Turtle Hatcheries* (Annex 1) was prepared by Dr Andrea Phillott in response to the needs in the Northern Indian Ocean sub-region. She further submitted *Best Practices In Sea Turtle Hatchery Management For South Asia*<sup>1</sup> (Annex 2) and a helpful infographic (Annex 3).
7. The 8<sup>th</sup> Meeting of the Advisory Committee recommends that further work on this subject is done, taking into account the needs in all the sub-regions. A corresponding measure has been added to Revision 2 of the Draft Work Programme.

---

<sup>1</sup> Phillott A.D. & Shanker K. 2018. Best practices in sea turtle hatchery management for South Asia. Indian Ocean Turtle Newsletter 27: 31-34.

## Best Practices for Sea Turtle Hatcheries

Dr. Andrea D. Phillott

Professor of Environmental Studies, FLAME University, Pune, Maharashtra, India

Senior Adjunct Fellow, Dakshin Foundation, Bengaluru, Karnataka, India

### Overview

1. Wherever possible, sea turtle eggs should be left to incubate in their original location on the beach.
2. When an unsustainable proportion of nests annually (dependent on species and population) are threatened by predation, poaching, tidal inundation, beach erosion etc, eggs may be moved to a hatchery- a protected, secure location on the nesting beach or nearby.
3. If eggs must be moved to a hatchery for protection, hatchling production can be optimised by use of best practices.
4. Table 1 in Phillott & Shanker (2018)<sup>1</sup> summarises best practices for:
  - Methods of egg collection, handling and transport
  - Incubation of eggs (nest depth and density)
  - Handling and release of hatchlings
  - Monitoring and evaluation of hatchling production
  - Record keeping
  - Training of staff and volunteers
5. Hatchery practices should be periodically assessed, and staff and volunteers should undergo regular training to ensure best practices are being implemented.
6. Threats to eggs at the natural beach should be periodically assessed to determine if eggs still need to be relocated to a hatchery.
7. Sea turtle eggs in their original location can be protected against poaching by community rangers or nest protectors, and against predators by barricades, cages, or nets.

---

<sup>1</sup> Phillott A.D. & Shanker K. 2018. Best practices in sea turtle hatchery management for South Asia. Indian Ocean Turtle Newsletter 27: 31-34.

# BEST PRACTICES IN SEA TURTLE HATCHERY MANAGEMENT FOR SOUTH ASIA

ANDREA D. PHILLOTT<sup>1#</sup> & KARTIK SHANKER<sup>2,3</sup>

<sup>1</sup>FLAME University, Pune, Maharashtra, India

<sup>2</sup>Centre for Ecological Sciences, Indian Institute of Science, Bengaluru, India

<sup>3</sup>Dakshin Foundation, Bengaluru, India

<sup>#</sup>andrea.phillott@flame.edu.in

Papers in Issue 27 of the *Indian Ocean Turtle Newsletter* highlight the extent to which countries in South Asia use hatcheries as an *ex situ* conservation strategy. Hatcheries are often perceived as 'safe' places to incubate sea turtle nests. Depending on the hatchery location and construction, eggs may be protected from predators, poachers, tidal inundation, and beach erosion. Hatcheries may also be used to raise community awareness about the biology and need for conservation of sea turtles, and provide opportunities for community revenue through ecotourism.

However, there should be ongoing threats to eggs in the natural environment for nests to be moved to a hatchery, as there are disadvantages to this practice. Hatching success in hatchery nests may be far lower than that of natural nests on the beach if poor hatchery management practices are employed, and sex ratios of hatchlings can be skewed if the nest temperatures within a hatchery differ from those on the natural beach. Operating a hatchery following best practice guidelines requires trained personnel and sufficient resources to protect and monitor nests throughout the nesting season (Mortimer, 1999; Shanker *et al.*, 2003).

Considering the disadvantages of incubating eggs in a hatchery, sea turtle nests should preferably remain *in situ* for the duration of the incubation period. Nests can be protected in their original location, for example, caged (Kurz *et al.*, 2011) or meshed (e.g. O'Connor *et al.*, 2017) to exclude predators, or relocated to a higher site on the beach if laid close to the high tide line (e.g. Tuttle & Rostal, 2010). Community awareness and conservation initiatives have proved successful at reducing the loss of nests to poachers (e.g. Rajakaruna *et al.*, 2009; West, 2010),

However, if moving eggs to a hatchery will ensure a much higher hatching success than the strategies described above, then hatchery location and construction,

methods of egg collection and transport, hatchery nest characteristics and density, and hatchling handling and release, should aim to maximise the number of hatchlings produced. Manuals relevant to the conservation of sea turtles in countries within the Indian Ocean and South East Asia (and globally) can be used as reference material for hatchery operations- Eckert *et al.* (1999), Ahmad *et al.* (2004) and STOI (2011). A visual summary of best practise for sea turtle hatchery management is available on the website *Sea Turtles of India* (<https://www.seaturtlesofindia.org/library/outreach-material/>). For quick reference, the best practices in hatchery management have been summarised in Table 1; references to studies that support the recommendations have also been provided.

We also recommend that hatchery managers periodically review the need for nests to be moved from their natural location, as changes in predator density and poaching activities may occur over time, removing the need for hatcheries to protect nests. Nests demonstrating a high hatching success and with a low risk of egg loss from depredation, poaching, tidal inundation, erosion, microbial invasion, etc should remain where they are laid.

## Literature cited:

Ahmad, A., T. Zulkifli, M.I. Mahyam, A.R. Solahuddin & Z. Nor Azman. 2004. *A Guide to Set-Up and Manage Sea Turtle Hatcheries in the Southeast Asian Region*. Marine Fisheries Resources Development and Management Department and Southeast Asian Fisheries Development Center (SEAFDEC).

Harry, J.L. & C.J. Limpus. 1989. Low-temperature protection of marine turtle eggs during long-distance relocation. *Australian Wildlife Research* 16: 317-320.

Limpus, C.J., V. Baker & J.D. Miller. 1979. Movement induced mortality of loggerhead eggs. *Herpetologica* 35: 335-338.

Maulany, R.I., D.T. Booth & G.S. Baxter. 2012. The effect of

Table 1. A summary of best management practices for sea turtle hatcheries.

	Best Practice Supporting Literature	Supporting Literature
Hatchery location	<ul style="list-style-type: none"> <li>• Choose a location at least one vertical metre above the highest high tide line and distant from tidal creeks, streams, river mouths etc to reduce risk of inundation or flooding and embryo mortality.</li> <li>• Minimise distance between nesting beach and hatchery to reduce transport time and potential for embryo mortality.</li> <li>• Provide a diversity of nest microhabitats (e.g. shade, slope) to reflect conditions on the nesting beach and avoid potentially skewing sex ratios of hatchlings.</li> <li>• Change location of hatchery every year to avoid accumulation of organic material and high microbial load. Change location of hatchery every year to avoid accumulation of organic material and high microbial load.</li> </ul>	Mortimer <i>et al.</i> (1999), Shanker <i>et al.</i> (2003), Ahmad <i>et al.</i> (2004); Spanier (2010); Maulany <i>et al.</i> (2012)
Hatchery construction (incl shading)	<ul style="list-style-type: none"> <li>• Enclose hatchery in fence constructed of chain link, wire mesh, barbed wire, cane, bamboo or slats as available.</li> <li>• Reinforce the base of the fence with 1-2m of 0.5cm-1.0cm mesh, buried to 50cm deep to prevent entry of burrowing predators.</li> <li>• Minimise the risk of lethal nest temperatures, especially late in incubation, by partially shading the hatchery with a material such as shade cloth or coconut thatch, or shading individual nests with thatch baskets. Permanently shaded hatcheries should monitor their nest temperatures to reflect those of in situ nests, to avoid skewing hatchling sex ratios from natural.</li> </ul>	Mortimer <i>et al.</i> (1999), Shanker <i>et al.</i> (2003), Ahmad <i>et al.</i> (2004)
Egg collection	<ul style="list-style-type: none"> <li>• Minimise movement and use of lights until turtle begins laying, to minimise disturbance and the risk the turtle will return to the sea without nesting.</li> <li>• Catch eggs as they are being laid by hand or into a clean plastic bag (be careful not to disturb the turtle by touch or movement during oviposition), or mark location of nest with a length of rope or coloured tape reaching from eggs to beach surface and remove eggs once turtle has finished nesting. Do not probe for nests with a stick or other implement, to avoid destroying eggs.</li> </ul>	Mortimer <i>et al.</i> (1999), Shanker <i>et al.</i> (2003), Ahmad <i>et al.</i> (2004)
Egg handling and transport	<ul style="list-style-type: none"> <li>• Rebury eggs in hatchery within 2-3hr of oviposition to minimise embryo mortality.</li> <li>• Eggs to be transported short distances more than 2hr after oviposition should be removed from the nest without vertical or horizontal rotation and egg orientation indicated by marking the top of eggs with a soft pencil. Original orientation should be maintained during transport.</li> <li>• Transport of eggs for long distances and/or long travel times may require low-temperature or hypoxic environments to maintain embryo viability.</li> <li>• Transport eggs in rigid containers to minimise rolling and potential embryo mortality.</li> </ul>	Limpus <i>et al.</i> (1979); Parmenter (1980); Harry & Limpus (1989); Mortimer <i>et al.</i> (1999), Shanker <i>et al.</i> (2003), Ahmad <i>et al.</i> (2004); Williamson <i>et al.</i> (2017)
Reburial of nests	<ul style="list-style-type: none"> <li>• Rebury eggs in a location within the hatchery that mimics the natural nest environment.</li> <li>• Dig hatchery nest to the same depth as the natural nest.</li> <li>• Mimic the shape of natural sea turtle nests (usually flask shaped with a narrower neck than base).</li> <li>• Place eggs individually into the nest; do not 'pour' eggs from a bucket or bag.</li> <li>• Cover the eggs with moist sand removed during nest construction; do not expose eggs to dry sand as there is a risk of desiccation.</li> <li>• Incubate a single clutch in the same hatchery nest; do not split clutches between nests, or combine clutches from different nests.</li> </ul>	Mortimer <i>et al.</i> (1999), Shanker <i>et al.</i> (2003), Ahmad <i>et al.</i> (2004); van de Merwe <i>et al.</i> (2006); Rusli & Booth (2016)
Nest density	<ul style="list-style-type: none"> <li>• Maintain a density of 1 nest/m<sup>2</sup> to minimise the effects of adjacent nests on temperature and respiratory gas availability, and allow space for hatchery workers to move.</li> </ul>	Mortimer <i>et al.</i> (1999), Shanker <i>et al.</i> (2003), Ahmad <i>et al.</i> (2004); Maulany <i>et al.</i> (2012)

Nest enclosures	<ul style="list-style-type: none"> <li>• Protect nests from predators by constructing cylindrical nest enclosures of rigid material ~60cm in diameter. (Avoid rigid wire as it can injure hatchlings.) Bury 10cm into sand for burrowing predators and cover with mesh or net for aerial predators.</li> </ul>	Mortimer <i>et al.</i> (1999), Shanker <i>et al.</i> (2003), Ahmad <i>et al.</i> (2004)
Hatching release	<ul style="list-style-type: none"> <li>• Predict emergence date, often 45-55 days after oviposition. The characteristic 'caving in' of sand above the nest indicates emergence will usually begin within 2-3 days.</li> <li>• Check enclosures every 30-60mins from afternoon to dawn and at other times when hatchlings may emerge (e.g. on overcast days and after rain) around the predicted emergence date.</li> <li>• Release hatchlings as soon as possible to prevent exhaustion, desiccation, loss of vigour, possible injury, or death from predators.</li> <li>• Release hatchlings in groups if possible to improve survival probability; however, early emergers should not be held until more hatchlings emerge as this practice can result in loss of vigour.</li> <li>• Randomise clutch release sites hundreds of metres apart to avoid creating fish feeding stations off the beach, which increases the risk of predation.</li> <li>• Allow hatchlings to crawl from the dune across the beach and enter the ocean unassisted to facilitate imprinting.</li> <li>• Manage observers to ensure hatchlings are not injured or their progress to the sea impeded; parallel lines ~20m apart on the beach give a mark for people to stand behind while hatchlings crawl between lines.</li> <li>• Ensure artificial lights are shielded during hatchling emergence and release, and after they enter the ocean to minimise disorientation.</li> <li>• If hatchlings emerge in heat of day or immediate release is not possible, hold hatchlings in a soft, damp cloth or sack in cool, dark place. Do not hold hatchlings in water as they will enter their 'swim frenzy' period and deplete energy reserves needed for survival and dispersal.</li> </ul>	Mortimer <i>et al.</i> (1999), Wyneken (2000); Pilcher & Enderby, 2001; Shanker <i>et al.</i> (2003), Ahmad <i>et al.</i> (2004); van de Merwe <i>et al.</i> (2013)
Hatchery records	<ul style="list-style-type: none"> <li>• Number each nest in the hatchery and associate with a standard data record form.</li> <li>• Complete a data sheet or data book entry for each nest, including information such as date of oviposition, clutch size, date of emergence, number of hatchlings, and (if recorded) SCL, weight, and scalation pattern.</li> <li>• Collect data from emerged hatchlings restrained in enclosure e.g. emergence date, and hatchling straight carapace length (SCL), weight and scalation pattern if possible.</li> </ul>	Mortimer <i>et al.</i> (1999), Shanker <i>et al.</i> (2003), Ahmad <i>et al.</i> (2004)
Monitoring and evaluation	<ul style="list-style-type: none"> <li>• Calculate incubation period as number of days between oviposition and emergence.</li> <li>• Excavate nest 2-3 days after the majority of hatchlings have emerged and calculate: <ul style="list-style-type: none"> <li>a) Hatching Success= (Number of hatched eggs/Total number of eggs) x 100</li> <li>b) Emergence Success= (Number of naturally emerged hatchlings/Total number of eggs) x 100</li> </ul> </li> <li>• Monitor nest temperature and hatchling sex ratio from a statistically valid proportion of nests in hatchery and compare with data from the natural beach/es for your population of sea turtles.</li> </ul>	Mortimer <i>et al.</i> (1999), Schäuble <i>et al.</i> (2002); Shanker <i>et al.</i> (2003)
Education and awareness	<ul style="list-style-type: none"> <li>• Create educational materials (e.g. posters) for visitors to the hatchery to raise awareness about sea turtle biology and conservation.</li> <li>• Encourage students and local wildlife enthusiasts to volunteer at the hatchery if possible.</li> </ul>	Shanker <i>et al.</i> (2003)
Personnel	<ul style="list-style-type: none"> <li>• Train employees and volunteers in sea turtle biology, conservation, and hatchery management techniques.</li> <li>• Provide access to general articles and manuals about sea turtle biology and hatchery practices.</li> <li>• Run or co-manage the hatchery with the local community when possible.</li> <li>• Collaborate with other community programmes in same area or elsewhere on the coast when possible.</li> </ul>	Shanker <i>et al.</i> (2003)

- incubation temperature on hatchling quality in the olive ridley turtle, *Lepidochelys olivacea*, from AlasPurwo National Park, East Java, Indonesia: Implications for hatchery management. *Marine Biology* 159: 2651-2661.
- Mortimer, J.A. 1999. Reducing threats to eggs and hatchlings: Hatcheries. In: *Research and Management Techniques for the Conservation of Sea Turtles* (eds. Eckert, K.L., K.A. Bjorndal, F.A. Abreu-Grobois & M. Donnelly). Pp. 175-178. IUCN/SSC Marine Turtle Specialist Group Publication No. 4.
- O'Connor, J.M., C.J. Limpus, K.M. Hofmeister, B.L. Allen & S.E. Burnett. 2017. Anti-predator meshing may provide greater protection for sea turtle nests than predator removal. *PLoS ONE* 12: e0171831.
- Parmenter, C.J. 1980. Incubation of the eggs of the green sea turtle, *Chelonia mydas*, in Torres Strait, Australia: The effect of movement on hatchability. *Australian Wildlife Research* 7: 487-91.
- Pilcher, N.J. & S. Enderby. 2001. Effects of prolonged retention in hatcheries on green turtle (*Chelonia mydas*) hatchling swimming speed and survival. *Journal of Herpetology* 35: 633-663.
- Rajakaruna, R.S., D.M.Naveen, J. Dissanayake, E.M.L. Ekanayake & K.B. Ranawana. 2009. Sea turtle conservation in Sri Lanka: Assessment of knowledge, attitude and prevalence of consumptive use of turtle products among coastal communities. *Indian Ocean Turtle Newsletter* 10: 1-13.
- Rusli, M.U. & D.T. Booth. 2016. Bigger clutch sizes save offspring energy during nest escapes. *Behavioral Ecology and Sociobiology* 70: 607-616.
- Schäuble, C., K. Ibrahim, A.R. Kassim, M. Hamann & J. Whittier. 2002. Monitoring hatchery success- What's worthwhile? In: *Proceedings of the 22<sup>nd</sup> Annual Symposium on Sea Turtle Biology and Conservation*. (comp. Seminoff, J.A.). NOAA Technical Memorandum NMFS-SEFSC-503, Pp. 308.
- Sea Turtles of India (SToI). 2011. A Comprehensive Field Guide to Research, Monitoring and Conservation (comps. Shenoy, S., T. Berlie & K. Shanker). Dakshin Foundation, Bangalore and Madras Crocodile Bank Trust, Mamallapuram, India. Pp. 148.
- Shanker, K., B.C. Choudhury & H.V. Andrews. 2003. *Sea Turtle Conservation: Beach Management and Hatchery Programmes*. A GOI-UNDP Project Manual. Centre for Herpetology/Madras Crocodile Bank Trust, Mamallapuram, Tamil Nadu, India.
- Spanier, M.J. 2010. Beach erosion and nest site selection by the leatherback turtle *Dermochelys coriacea* (Testudines: Dermochelyidae) and implications for management practices at Playa Gandoca, Costa Rica. *Revista de Biología Tropical* 58: 1237-1246.
- Tuttle, J. & D. Rostal. 2010. Effects of nest relocation on nest temperature and embryonic development of loggerhead sea turtles (*Caretta caretta*). *Chelonian Conservation and Biology* 9: 1-7.
- van de Merwe, J., K. Ibrahim & J. Whittier. 2006. Effects of nest depth, shading, and metabolic heating on nest temperatures in sea turtle hatcheries. *Chelonian Conservation and Biology* 5: 210-215.
- van de Merwe, J.P., K. Ibrahim & J. M. Whittier. 2013. Post-emergence handling of green turtle hatchlings: Improving hatchery management worldwide. *Animal Conservation* 16: 316-323.
- West, L. 2010. A multi-stakeholder approach to the challenges of turtle conservation in the United Republic of Tanzania. *Indian Ocean Turtle Newsletter* 11: 44-50.
- Williamson S.A., R.G. Evans & R.D. Reina. 2017. When Is embryonic arrest broken in turtle eggs? *Physiological and Biochemical Zoology* 90: 523-532.
- Wyneken, J. 2000. The migratory behaviour of hatchling sea turtles beyond the beach. In: *Sea Turtles of the Indo-Pacific*. (eds. Pilcher, N.J. & G. Ismail). Pp. 121-142. ASEAN Academic Press, London.




# BEST PRACTICES SEA TURTLE HATCHERIES

Sea turtle nests may be relocated to a hatchery as part of a conservation, research or outreach programme. From a conservation perspective, nests should only be relocated to a hatchery if it is absolutely essential i.e. if there is a significant danger of predation, tidal flooding or other threats.



✓ The hatchery should be located about 20 to 30m from the high tide line to prevent flooding and should be close to the nesting site.




✓ The hatchery can be constructed with poles and mesh; nests should be at least 2 metres apart.



✓ Hatchlings should be released at night immediately after hatching and allowed to crawl down the beach into the sea.



✓ Close to the emergence date, each nest should be covered with a basket/plastic mesh.



✓ The shape of the nests should be similar to that of natural nests.



✓ Basic data should be recorded on a placard at the nest, and a detailed data book should be maintained.



✗ The hatchery should not be covered on top or in the shade of trees



✗ The hatchlings should never be held in buckets of water or sand.



✗ The hatchery should not be established at the same site for two consecutive years.



✗ Hatchlings should not be released at the same site everyday.

