



**MEMORANDUM OF UNDERSTANDING  
ON THE CONSERVATION OF  
MIGRATORY SHARKS**

CMS/Sharks/MOS2/Doc.8.2.3

18 September 2015

Original: English

---

Second Meeting of the Signatories  
San José, Costa Rica, 15-19 February 2016  
Agenda Item 8

**PROPOSAL FOR THE INCLUSION OF THE REEF MANTA RAY,  
*MANTA ALFREDI*, IN ANNEX 1 OF THE CMS MEMORANDUM OF  
UNDERSTANDING ON THE CONSERVATION OF MIGRATORY SHARKS**

*(Prepared by the Secretariat)*

1. The present proposal for the inclusion of the entire population of the Reef Manta Ray (*Manta alfredi*) in Annex 1 to the MOU represents the original proposal for inclusion of the species in CMS Appendix I and II, submitted as UNEP/CMS/COP11/Doc.24.1.9/Rev followed by an Addendum to provide additional information by the Government of Fiji to the 11th Meeting of the Conference of the Parties (CMS COP11). The proposal was subsequently adopted by the Parties.
2. As agreed at the 1<sup>st</sup> Meeting of the Signatories (MOS1) and in line with the procedure explained in CMS/Sharks/MOS2/Doc.8.2.1, the original proposal is now being resubmitted for consideration by the Second Meeting of the Signatories (MOS2). Signatories are requested to consider the inclusion of *Manta alfredi*, in Annex 1 of the Memorandum of Understanding on the Conservation of Migratory Sharks (Sharks MOU) based on the information provided in this document.
3. The Advisory Committee of the MOU has presented a review of the proposal in CMS/Sharks/MOS2/Doc.8.2.10 in which the Committee recommends the entire population of *Manta alfredi* for inclusion in Annex 1.



**PROPOSAL FOR INCLUSION OF SPECIES ON THE APPENDICES OF THE  
CONVENTION ON THE CONSERVATION OF MIGRATORY SPECIES OF  
WILD ANIMALS**

*(Originally submitted as UNEP/CMS/COP11/Doc.24.1.9/Rev.1 to CMS COP11 by the  
Government of Fiji on 4 November 2014)*

- A. PROPOSAL:** Inclusion of the species *Manta alfredi* (Reef Manta Ray), Genus *Manta*, Subfamily *Mobulinae*, in Appendix I and II
- B. PROPONENT:** Government of the Fiji Islands
- C. SUPPORTING STATEMENT:**
- 1. Taxon**
- 1.1 Class:** Chondrichthyes, subclass Elasmobranchii
- 1.2 Order:** Rajiformes
- 1.3 Subfamily:** Mobulinae
- 1.4 Genus and species:** Genus *Manta* (Bancroft 1829): *Manta alfredi* (Krefft 1868) and any other putative *Manta* species.
- Scientific Synonyms:** *Deratoptera alfredi* (Krefft 1868)  
*Manta fowleri* (Whitley 1936)
- 1.5 Common Names:** English: Reef Manta Ray, Prince Alfred's Ray, Inshore Manta Ray, Coastal Manta Ray, Resident Manta Ray

**Overview**

- i. *Manta alfredi*, a globally distributed and highly migratory species, is proposed here for listing on CMS Appendix I and II. This iconic and highly vulnerable species would benefit from strict range state protections under a CMS Appendix I listing as well as collaborative management initiated under a CMS Appendix II listing, since it is a slow reproducing, commercially exploited aquatic species that is in decline. In addition, international cooperation under the Appendix II listing would be greatly facilitated by adding all species of the Subfamily Mobulinae (Genus *Manta* and Genus *Mobula*) to Annex I of the CMS Sharks MoU. Increasing international trade in Mobulinae gill plates, and to a lesser degree skins and cartilage, and unregulated bycatch in industrial and artisanal fisheries, have led to significant rates of decline in population sizes in recent years.

App I listing would encourage range state Parties where *M. alfredi* are targeted to seek help with implementing measures to protect this species and enable artisanal fishers to benefit from the much more lucrative and sustainable tourism revenues this species generates. For example in Mozambique, *Manta alfredi* and the closely related *Manta birostris* (already included on Appendix I) contribute over US\$13 million annually to tourism revenues. However, off Praia do Tofo, an important manta and whale shark tourism area in southern Mozambique, artisanal fishers opportunistically target mantas for their low value meat. In this area, Rohner et al. (2013) observed a decline in *Manta alfredi* abundance of 88% over

only 8 years (less than a third of one generation period for this species).

In addition, new evidence of growing demand for manta and mobula gill plates and expansion of this trade threatens to motivate the emergence of new, targeted manta fisheries in range states where *M. alfredi* are currently not protected. CMS parties that are *M. alfredi* range states, which currently do not have protection measures in place for *M. alfredi*, include Australia, Comoros-Mayotte (France) Cook Islands, Djibouti, Egypt, Fiji, India, Madagascar, Mozambique, New Caledonia (France), Pakistan, Palau, Philippines, Saudi Arabia, Seychelles, South Africa, and Yemen. The CMS Appendix I listing, therefore, will help this species by curbing existing targeted fisheries and preventing emergence of new fisheries in response to demand from the mobulid gill plate trade.

- ii. *M. alfredi* are slow-growing, large-bodied migratory animals with small, highly fragmented populations that are sparsely distributed across the tropical and oceans of the world. They have among the lowest fecundity of all elasmobranchs, typically giving birth to a single pup with a gestation period of approximately one year, placing them into FAO's lowest productivity category. Global population numbers are unknown, but thought to be declining across their range. Their biological and behavioural characteristics (low reproductive rates, late maturity and aggregating behaviour) make these species particularly vulnerable to over-exploitation in fisheries and extremely slow to recover from depletion.
- iii. Populations of *M. alfredi* have not been assessed but appear to be generally small, sparsely distributed, and characterized by very low productivity and aggregating behaviour, leaving them especially vulnerable to exploitation and with limited capacity to recover from depletion. Additional impacts from the overexploitation of manta rays include the significant economic consequences to sustainable ecotourism operations estimated to generate US\$140 million annually due to the presence of manta rays (O'Malley *et al.* 2013). Non-consumptive utilization of manta rays through tourism can yield much larger and longer-term benefits to range states than manta ray fisheries (Anderson *et al.* 2010, Heinrichs *et al.* 2011, O'Malley *et al.* 2013), which are not likely to be sustainable even at moderate levels (Dulvy *et al.* 2014, Rohner *et al.* 2013).
- iv. The prebranchial appendages (or gill plates), which *M. alfredi* use to filter planktonic food from the water, are highly valued in international trade. Cartilage and skins are also traded internationally. A single mature *M. alfredi* can yield up to 5 kilos of dried gills that retail for up to US\$390 per kilo in China. Since there are no specific import-export codes for *Manta spp.* gill plates and trade records for cartilage and skins are generally not species-specific, international trade levels, patterns and trends cannot be accurately documented. Existing DNA tests and visual ID guides can enable informed non-experts to distinguish *Manta spp.* and their parts and derivatives in trade from other species.
- v. *M. alfredi* are caught in commercial and artisanal fisheries throughout their global warm water range in the Atlantic, Pacific and Indian Oceans. Directed fisheries primarily utilize harpoons and nets, while significant bycatch may occur in purse seine, gill and trawl net fisheries targeting other species. The high value of gill plates has driven increased target fishing pressure for all *Manta spp.* in key range states, with the largest landings observed in Indonesia, Sri Lanka and Mozambique. Fisheries in other countries may also be significant, but landings data from most locations are not readily available. The recent

increase in demand for gill plates has resulted in dramatic increases in fishing pressure, with many former bycatch fisheries having become directed commercial export fisheries. There are also recent reports of mantas being ‘gilled’ (gills removed and the carcasses discarded) at sea (D. Fernando pers. obs.).

- vi. There have been no stock assessments, official monitoring, catch limits or management of *M. alfredi* fisheries in the waters of range States with the largest fisheries. Regional Fishery Management Organizations (RFMOs) have not taken any measures to minimize high seas bycatch of *Manta spp.* Incidental landings and discards are rarely recorded at the species level. *M. alfredi* are legally protected in a few countries and in some small Marine Protected Areas, but most laws protecting manta rays define “Manta Ray” as “*Manta birostris*”, leaving *M. alfredi* unprotected.
- vii. While there are no historical baseline population data, recent declines have been reported in key *M. alfredi* range states, including Indonesia and Mozambique (M. Erdmann, pers. comm., S. Lewis, pers. comm., J. Friedlander, pers. comm., Rohner et al. 2013).
- viii. Following consideration of a taxonomic review prepared by the IUCN SSC Shark Specialist Group (Fowler & Valenti/SSG 2007), the CMS Scientific Council agreed in March 2007 (CMS SCC14) that these threatened migratory species meet the criteria for listing on the Appendices and should be considered by the Conference of Parties to CMS. *Manta birostris*, the other species in the genus *Manta*, was added to CMS Appendix I and II at the 10th CoP in 2011. Until the recent split of the genus *Manta* (Marshall et al. 2009, Kashiwagi et al. 2012), all manta rays were classified as *M. birostris*; the two species share highly similar biological and behavioural characteristics and face the same threats.

## 2. Biological data

The subfamily Mobulinae encompasses two genera: *Manta* and *Mobula*. This group is characterized by the presence of one lobe on each side of the head, wing-like pectoral fins, terminal mouth and a stingless tail (Notarbartolo-Di-Sciara 1987a). All are planktivorous, feeding on zooplankton and (in the case of several of the *Mobula spp.*) small schooling fishes. The genus *Manta* was previously considered monotypic; a focused genetic study has confirmed that *M. birostris* and *M. alfredi* are two distinct species (Kashiwagi et al. 2012). Descriptions or photographs can be used to verify accounts to the species level.

*Manta spp.* are large-bodied, pelagic, planktivorous rays. *M. birostris* can grow to over 7 meters wingspan (disc width or DW; Marshall et al. 2009) with anecdotal reports up to 9 meters (Compagno 1999). *M. alfredi* grows to an average 3.5 meters DW, and a maximum of 5 meters DW (Marshall et al. 2011b). *Manta spp.* are distinguished by their large diamond-shaped body with elongated wing-like pectoral fins, ventrally placed gill slits, laterally placed eyes, wide terminal mouths, and paired cephalic lobes. Melanistic (black) and leucistic (white) colour morphs occur in both species (Marshall et al. 2009). Most *Manta spp.* show a counter-shading pattern (black dorsally and white ventrally) and have unique spot patterns on their ventral surface that do not change over time and help identify individuals (Clark 2001, Marshall et al. 2008, Kitchen-Wheeler 2010, Deakos et al. 2011).

Mantas are slow-growing and long-lived with low fecundity and reproductive output and long generation times (estimated at 25 years). Longevity is estimated to be at least 40 years (Marshall et al. 2011b,c) and natural mortality is thought to be low (Couturier et al. 2012).

Mantas are among the least fecund of all elasmobranchs (Couturier *et al.* 2012; Dulvy *et al.* 2014), bearing only one pup on average every two to three years (Marshall *et al.* 2011a,b, with a gestation period of 10– 14 months (Homma *et al.* 1999; Marshall *et al.* 2009; M. de Rosemont pers. comm.) and reaching maturity at ~10 years (Marshall *et al.* 2011b,c). Earlier age at maturity (~3-6 years) was estimated in males in one subpopulation in Kona, Hawaii (Clark 2010). Later maturity (15 years or more) and lower reproductive rates (one pup every seven years) have been observed for female *M. alfredi* in a subpopulation in the Maldives (G. Stevens in prep.). With such conservative life history characteristics, a female manta ray can produce no more than 4-15 pups over her lifetime. Subpopulations are therefore exceptionally vulnerable to extirpation, slow to recover once depleted; the possibility of successful re-colonization is low. In local populations with low anthropogenic influences, *Manta alfredi* females have been found to need to produce at least one pup for every 3.1 years on average to achieve a small, but positive population growth (Kashiwagi 2014).

## 2.1 Distribution and range states (current and historical)

*Manta* spp. are circumglobal in range (see Annex I), with the two described species sympatric in some locations and allopatric in others (Kashiwagi *et al.* 2011). *M. alfredi* is found in tropical and subtropical waters (Marshall *et al.* 2009, Kashiwagi *et al.* 2011, Couturier *et al.* 2012). *Manta alfredi* populations are sparsely distributed and highly fragmented, likely due to their resource and habitat needs.

## 2.2 Population (estimates and tendencies)

In extensively studied *M. alfredi* subpopulations in Mozambique, eastern Australia and the Maldives, a significant female bias has been observed, with the majority in Mozambique considered to be mature (Marshall *et al.* 2011a, Couturier *et al.* 2014; G. Stevens, unpubl. data). In an *M. alfredi* subpopulation in Maui, Hawaii, the sex ratio is close to parity with juveniles and adults present. This study also suggests that juveniles may segregate from the adult population, residing in areas where they are less vulnerable to predation (Deakos *et al.* 2011). In Ningaloo, Australia, the distribution of males to females and adults to juveniles fluctuates throughout the year, but mature females consistently dominate (McGregor 2009). Of three *M. alfredi* aggregation sites surveyed in eastern Australia, only the largest site exhibited a significant female bias while the other two showed no bias (Couturier *et al.* 2011).

Subpopulations of the *M. alfredi* appear, in most cases, to be small (less than 1,000 individuals). Photo-identification studies at aggregation sites in Hawaii (Deakos *et al.* 2011), the Yaeyama Islands Japan (Kashiwagi 2014), southern Mozambique (Marshall *et al.* 2011), and the east coast of Australia (Couturier *et al.* 2014) have produced sighting records or estimated annual or sub-population sizes of approximately 100 to 700 individuals, despite some being active for many decades. The one exception is the Maldives with 3,300 individuals identified throughout the 26 atolls that make up the archipelago (G. Stevens, unpublished data 2012) and population estimates for the entire archipelago reaching 9,677 individuals (Kitchen-Wheeler *et al.* 2011). These preliminary studies at major aggregation sites suggest resident population sizes of *Manta alfredi* are generally small, with some areas having large seasonal influxes.

Population reductions of *Manta alfredi* have been reported in areas with active fisheries or significant by-catch such as southern Mozambique where an 88% decline in observation sighting records has occurred over a nine year period (Rohner *et al.* 2013), well under one

generation estimated at 25 years for *Manta* spp. (Marshall et al. 2011b,c). Globally a decline of 30% is strongly suspected (Marshall et al. 2011b,c) and currently *M. alfredi* is listed as Vulnerable on the IUCN Red List of Threatened Species with a decreasing population trend.

### 2.3 Habitat (brief description and tendencies)

*M. alfredi* are more commonly sighted inshore, but are also observed around offshore coral reefs, rocky reefs and seamounts. This species is often resident in or along productive nearshore environments, such as island groups, atolls, or continental coastlines, and may also be associated with areas or events of high primary productivity (e.g., upwelling) (Homma et al. 1999, Dewar et al. 2008, Kitchen-Wheeler 2010, Anderson et al. 2011, Deakos et al. 2011, Marshall et al. 2011b).

### 2.4 Migration (types of movement, distances, proportion of the population that migrates)

Telemetry studies and photographic matching studies show that *M. alfredi* is highly mobile, and while they typically exhibit smaller home range sizes than *M. birostris*, this species often undertakes regular or seasonal migrations (up to several hundred kilometers) most likely to exploit productive feeding grounds (Anderson et al. 2011, Jaine et al. 2012)

Long-term sighting records of *M. alfredi* at established aggregation sites suggest that this species, is quite resident to tropical water habitats, exhibits smaller home ranges, and shows philopatric behaviour for specific critical habitats like feeding areas and inshore reefs that support cleaning stations (Kashiwagi 2014, Kitchen-Wheeler et al. 2011, Marshall et al. 2011a, , Deakos et al. 2011, Clark 2010, Couturier et al. 2014).

Telemetry studies and broad-scale photo-matching studies are increasingly showing that *M. alfredi* is capable of traveling significant distances in short periods of time. Couturier et al. 2014 showed *Manta alfredi* undertaking migrations of up to 650 kilometers in a 6-month period along the eastern coast of Australia. Germanov and Marshall 2014 showed *Manta alfredi* making regular migrations between the Komodo National Park and the manta ray sanctuary of Nusa Penida in Indonesia (distances of up to 450km) in as little as 33 days. A study by Duinkerken 2010 showed *M. alfredi* in southern Mozambique traveling between sites up to 95 kilometers apart in as little as 40-hours, with maximum rate of movements of 3.7 km h<sup>-1</sup> suggesting that this species is capable of traveling large distances in short periods of time. In most cases these longer range migrations occurred along continuous continental coastlines rather than across island chains or large bodies of water. However, a recent study by Jaine et al. 2014 using satellite telemetry off eastern Australia found *Manta alfredi* traveling up to 155km offshore to feed, swimming up to 2,441 km (not a straight-line distance) in 118 days. Braun et al. 2014 also found satellite tagged *M. alfredi* to be using offshore environments up to 28% of their monitored track.

While some sub-populations of *M. alfredi* may be deterred by physical barriers, like open expanses of sea (Deakos et al. 2011) it is clear that in certain circumstances or in some locations *M. alfredi* may undertake regular longer distance movements. While no international migrations have been documented in the literature, the distances that *M. alfredi* have been demonstrated to regularly undertake would suggest that they are capable and are most likely moving through and using habitats in adjoining countries in parts of their distribution. Photo matches of the same individuals using habitats in Inhambane Mozambique and those in Ponta de Oro (less than 1.5 km from the boarder of South Africa) are illustrations of this point

(Manta Matcher public records-

<http://mantamatcher.org/individuals.jsp?langCode=en&number=MZ0803A>).

Daily diurnal migrations are reported in *M. alfredi* with individuals using inshore environments like shallow reef cleaning stations and coastal feeding grounds during daylight hours and deeper water/off shore habitats in the evening hours (Dewar et al. 2008, Marshall 2009, Anderson et al. 2011, Duinkerken 2010, Braun et al. 2014). Migrations from areas of protection to offshore environments (Braun et al. 2014, Jaine et al. 2014), or through areas fraught with anthropogenic threats (Germanov and Marshall 2014) could put *M. alfredi* at risk, even if their inshore or main critical habitats are protected. In other more closed subpopulations with little to no exchange between neighboring sub-populations (Deakos 2012, Kashiwagi 2014) unsustainable fishing or anthropogenic influences could deplete a single population quite rapidly, with little chance of recovery or re-population over time.

### **3. Threat data**

The greatest threat to *M. alfredi* is unmonitored and unregulated directed fisheries, increasingly driven by the escalating international trade demand for their gill plates (Heinrichs et al. 2011, Whitcraft et al. 2014). The gill rakers are predominately used in Asian health tonics, purported to treat a wide variety of conditions. Artisanal fisheries also target reef manta rays for local consumption (B. Newton, pers. comm., J. Hartup, pers. comm., Rohner et al. 2013).

Manta species in general are easy to target because of their large size, slow swimming speed, aggregating behaviour, predictable habitat use, and lack of human avoidance. They are killed or captured by a variety of methods including harpooning, longlining, netting and trawling (Couturier et al. 2012, White et al. 2006, Heinrichs et al. 2011, Fernando and Stevens 2011). Targeting of these rays at critical habitats or aggregation sites, where individuals can be caught in large numbers in a short time frame, is a potentially serious threat (Couturier et al. 2012). Their conservative life history also constrains their ability to recover from a depleted state and they are not likely to be able to tolerate high catch levels, given their very low reproductive potential (Dulvy et al. 2014).

Industrial fisheries or inshore gill net fisheries, which are by nature quite indiscriminate, can result in manta ray by-catch. This species also faces many anthropogenic threats, particularly entanglement (in phantom nets, mooring lines, anchor lines and fishing lines), boat strikes and sport fishing-related injuries. Additional threats include habitat destruction, pollution, climate change, oil spills and ingestion of marine debris such as micro plastics (Couturier et al. 2012).

#### **3.1 Direct threats to the population (factors and intensity)**

Historically, subsistence fishing for Manta spp. occurred in isolated locations with simple gear, which limited the area and time fishermen could hunt. In recent years, however, fishers have begun targeting Manta spp. with modern fishing gear and expanding their fishing range and season, primarily in response to the emerging market for dried mobulid gill plates (Dewar 2002, White et al. 2006, Rajapackiam et al. 2007, White and Kyne 2010, Heinrichs et al. 2011, Setiasih et al. in prep., Fernando and Stevens 2011). This increase in fishing pressure is driving regional Manta spp. subpopulations toward commercial extinction (Dewar 2002, White et al. 2006, Heinrichs et al. 2011).

As *M. alfredi* was only recognized a separate species from the other manta species, *M. birostris*, in 2009 (Marshall et al., 2009, Kashiwagi et al. 2012), data prior to this does not distinguish among the two. However their relatively smaller home range, tendency for nearshore residency and philopatric behaviour to critical habitats such as inshore reefs makes *M. alfredi* a high target for fisheries, and it is likely they represent a significant proportion of the reported manta catch. Today, the largest documented manta fishing and exporting Range States are Indonesia (Indonesia introduced complete protective legislation for both *Manta* spp. within its territorial waters in March 2014), Sri Lanka, and India. However, gill plate vendors in China also report Malaysia, Vietnam, South Africa, South America, the Middle East and the South China Sea as source regions for mobulid gill plates (Whitcraft et al. 2014), and high international trade demand is likely to stimulate directed and opportunistic fisheries elsewhere. In light of the lack of published fishery data at the species level, the very recent separation of the genus into two species (Marshall et al. 2009, Kashiwagi et al. 2012), the species' extremely high biological vulnerability (Dulvy et al. 2014) and the rapidly escalating trade demand for manta ray gill plates - regardless of species (Whitcraft et al. 2014), the precautionary approach is recommended to prevent the spread of unsustainable fisheries and population depletions.

**Pacific Ocean:** Opportunistic hunting of a small *M. alfredi* population has recently been reported in the islands of Tonga (B. Newton, pers. comm.) and Micronesia (J. Hartup, pers. comm.). Because of their isolation and low numbers, such local subpopulations of *M. alfredi* are extremely vulnerable to any fishing pressure. Neither of the *Manta* species are being targeted for commercial fishing or subsistence harvest across Fijian waters, and thus now are being currently protected under the Endangered and Protected Species Act, and its attending regulation. All species of *Manta* from September 2014 will be protected under CITES Appendix II.

**Indo-Pacific:** *Manta* spp. fisheries have been observed in Indonesia in Lamakera and Lamalera (Nusa Tenggara), Tanjung Luar (Lombok), Cilacap (Central Java) and Kedonganan (Bali) (Dewar 2002, White et al. 2006, Barnes 2005). Most fisheries are targeted and have arisen or greatly increased over the past ten years. In and around the Wayag and Sayang Islands in Raja Ampat, Indonesia, where shark populations have collapsed, shark fishermen have reportedly begun to target *Manta* spp. (Donnelly et al. 2003). In Lamakera, when motorized boats replaced traditional dugout canoes to target *Manta* spp., catch rates increased by an order of magnitude above historic levels (Dewar 2002). Lamakera fishermen reported in 2002 that previously manta rays had occurred in the channel near the village, but were no longer seen nearshore, suggesting possible local extirpation of a reef manta population (Dewar 2002).

While peer reviewed sources have only reported landings of *M. birostris* in these locations, these sources pre-date the separation of the genus *Manta* from one species (*M. birostris*) to two species (*M. birostris* and *M. alfredi*). The proportion of each species in the landing is therefore unclear. Recent reports from Tanjung Luar, Lombok, however, indicate increased pressure on local *M. alfredi* populations (P. Hilton, pers. comm.). Dive operators and tourists have also reported seeing manta rays in fish markets close to known *M. alfredi* aggregation sites in Sangalaki, Borneo (E. Oberhauser, pers. comm.) and the Ende market in Flores near Komodo National Park (B. Pilkington-Vincett, pers. comm.).

There is anecdotal evidence of manta population declines and possible extirpations at three additional sites in Indonesia; Pulau Banyak, Lembeh Strait and Alor Island. Patrol staff at

Yayasan Pulau Banyak off the west coast of Sumatra report that local fishermen catch manta rays as by-catch in gill nets and that manta sightings have become much less frequent, suggesting a possible population decline as a result of by-catch fishing pressure (S. Lewis, pers. comm.). In 1997 in the Lembah Strait region of North Sulawesi, 1,424 manta rays were caught in large trap nets set in a migratory channel designed to catch pelagic fish and marine mammals entering the mouth of the Lembah Strait, just off the Tangkoko Nature Reserve (Anon 1997). Use of these nets was prohibited, but they were found to be in use again shortly afterwards (White et al 2006). Following a legal battle between the Indonesian Ministry of Environment and the Taiwanese company operating the trap net, the nets were permanently removed in 1998. Prior to deployment of the trap nets, Lembah dive operators reported the presence of a resident manta ray population, and divers regularly snorkeled with mantas feeding in the channel (J. Friedlander, pers. comm.). However mantas have not been seen in the channel since this time (M. Erdmann, pers. comm.). Local Alor fishermen reported that approximately ten years ago off the west coast of Alor Island, in the channel between Alor and Pantar Islands, a local village started installing drift nets in the middle of the channel targeting mackerel. The nets were reportedly 50m wide and set at a depth of 18-20m. Manta rays were caught as by-catch in the nets and within five years mantas were no longer seen in the region, suggesting that the nets may have caused a population extirpation. Prior to the installation of these nets manta rays were reportedly seen commonly, usually at the surface in groups of two but sometimes in larger groups of 10-15 individuals. Despite a lack of photographic evidence, local fishermen reported that the mantas were approximately 3m from wing tip to wing tip thus suggesting that they were likely to have been *M. alfredi*. (M. Erdmann, pers. comm, S. Lewis, pers. comm.).

**Indian Ocean:** Targeted fisheries are reported Thailand (R. Parker, pers. comm.), the Philippines (Alava et al. 2002 – now legally prohibited), and several locations in Africa, including Tanzania and Mozambique, where annual landings of ~35-50 *M. alfredi* are reported from less than 5% of the coastline, (Marshall and Bennett 2010).

### 3.2 Habitat destruction (quality of the changes, quantity lost)

The loss of some coral reef habitats, which provide food, cleaning stations and reproductive areas, could have a negative impact on *Manta* spp. (Deakos 2010). Alterations to terrestrial ecosystems have also been shown to affect *Manta* spp. populations. At Palmyra Atoll in the Pacific, a study linked declines in the manta rays' planktonic food source to areas where native trees have been replaced by human propagated palms, revealing a complex interaction chain linking trees to manta rays (McCauley et al. 2012). *Manta* spp. are also likely to be susceptible to oil spills and pollution because of their wide-ranging near-shore habitat preferences (Notarbartolo di Sciara 2005, Handwerk 2010).

Chin and Kyne (2007) estimated that mobulid rays (Genus *Manta*; Genus *Mobula*) are the pelagic species most vulnerable to climate change, since plankton, a primary food source, may be adversely affected by the disruption of ecological processes brought about by changing sea temperatures. In the Republic of Maldives, over a three year period (2009-2012), despite intensive directed research, there were no recorded pregnancies amongst a subpopulation of over 659 individually identified mature female *M. alfredi* (G. Stevens in prep). This scarcity of pregnancies correlates directly with un-seasonally weak monsoonal winds in the region, which should drive the nutrient upwellings that lead to the rich

productivity of the Archipelago upon which the manta ray directly depend (Anderson et al. 2011, G. Stevens pers. comm.). These broad scale fluctuations in the productivity of the Maldivian waters are reflected in catch rates of the local tuna fishery, which have been linked to wider climatic patterns such as the El Niño Southern Oscillation (ENSO) (Anderson 1999). Other habitat threats that affect *Manta* spp. populations include marine debris such as, ghost nets and plastics, and pollution from vessels.

### 3.3 Indirect threats

*Manta* spp. are a bycatch of myriad fisheries targeting other species throughout the Atlantic, Pacific, and Indian Oceans, but are most frequently bycaught in purse seines, gillnets, and longlines (all commonly used in tuna fisheries). Bycatch data are collected in only a few fisheries and, when they are, *Manta* spp. are often recorded under various broad categories such as “Other”, “Rays”, or “Batoids”, with a breakdown by species almost never recorded (Lack and Sant 2009, Camhi *et al.*).

Numbers of animals released alive are only rarely recorded, while visual identification field guides for *Manta* and *Mobula* spp. have only recently been published (G. Stevens, 2011). As such, *Manta* spp. have generally been overlooked in most oceanic fisheries reports, with very little effort to properly identify or accurately record the species caught (Chavance *et al.*, 2011, G. Stevens, pers. comm.).

*M. alfredi* are also threatened by entanglement (in phantom nets, mooring lines, anchor lines and fishing lines), boat strikes and sport fishing-related injuries. Additional threats include habitat destruction, pollution, climate change, oil spills and ingestion of marine debris such as micro plastics (Couturier *et al.* 2012).

### 3.4 Threats connected especially with migrations

See new additions to migrations section

### 3.5 National and international utilization

All utilisation and trade in the products of *Manta* spp. is derived from wild-caught animals. Records cannot be quantified fully, due to a lack of species and product-specific codes, catch, landings, and trade data. All available information, however, indicates that fisheries are trending from bycatch to more targeted operations primarily to supply gill plates to Asian markets (Fernando and Stevens in prep, Heinrichs et al. 2011, Setiasih et al. in prep., Dewar 2002, Marshall et al. 2011b,c). For example, fishermen in Sri Lanka used to avoid setting their nets where *Manta* spp. were known to occur, and any rays caught incidentally were released, often alive, at sea. Following the rapid growth of the gill plate trade over the past decade, however, fishermen now land all *Manta* spp. and have recently begun removing the gill plates at sea, discarding the remaining low-value carcass (D. Fernando, pers. comm.).

The ~35-50 *M. alfredi* are taken annually in southern Mozambique (Marshall and Bennett 2010) are used domestically for consumption although their meat is not considered valuable or even the preferred diet of local fishing communities. There is no documented domestic use of *Manta* spp. gill plates in the three largest *Manta* spp. fishing range States (Indonesia, Sri Lanka and India) (Heinrichs et al. 2011, Fernando and Stevens in prep, Setiasih et al. in prep.). The relatively low-value meat of *Manta* spp. taken in these and other domestic

fisheries is used locally for shark bait, animal feed, and human consumption or discarded, while high value products (primarily gill plates, also skin and cartilage) are exported for processing elsewhere (Heinrichs et al. 2011, Setiasih et al. in prep., Fernando and Stevens in prep, Marshall et al. 2011b,c, Booda 1984, C. Anderson, pers. comm., D. Fernando pers. comm.).

Small numbers of *M. birostris* and *M. alfredi* are also caught and transported to aquariums for use in large display tanks in the US, Bahamas, Portugal, Japan, Singapore and South Africa. Uchida (1994) reported the number of surviving days for manta rays in captivity from 1 to 1,943.

An analysis of Manta tourism relative to fisheries value in Indonesia, home to the largest fishery for Manta spp. until protective legislation came into effect in March 2014, estimated tourism revenues in excess of USD 15 million per year compared with fishery revenues of ~USD 442 thousand annually (O'Malley et al. 2013). Dive tourism in Yap is focused almost exclusively on Manta ray encounters, with an annual value estimated at USD 4 million (B. Acker, unpubl.). Tourism operations focused on viewing marine megafauna such as manta rays bring millions of dollars in revenue annually primarily to local communities (Norman and Caitlin 2007, Pine et al. 2007, Brunnschweiler 2009, Tibirica et al. 2009, Jones et al. 2009, Graham 2004, Martin and Hakeem 2006, Hara et al. 2003, Topelko and Dearden 2005). In the Maldives, for example, direct revenue from manta dive and snorkel excursions was estimated to generate over USD 8.1 million per year during 2006–2008 (Anderson et al. 2010).

Globally, the direct economic impact of Manta spp. dive tourism is estimated at USD 140 million per year (O'Malley et al. 2013). Meanwhile, tourism opportunities in a number of range States have still not been explored. These existing and potential tourism revenues are significantly greater than the estimated value of USD 5 million per year for the global Manta spp. gill plate trade (Heinrichs et al. 2011). The development of high value community-based whale shark tourism in the former fishing range States of the Philippines and Indonesia illustrates the potential for Manta tourism to provide long-term, sustainable income to many coastal communities, if short-term boom and bust fisheries are avoided.

## **4. Protection status and needs**

### **4.1 National protection status**

While the capturing and killing of manta rays is banned in several countries, most of these laws define “Manta ray” as “*M. birostris*”, leaving few legal protections for *M. alfredi*. Currently only Indonesia, the state of Yap (Federated States of Micronesia), the U.S. states / territories of Hawaii, Florida, Guam and the Commonwealth of the Northern Mariana Islands, and the Australian Indian Ocean territories Christmas Island and Cocos Keeling Islands have laws protecting both Manta species. Yap’s Manta Ray Sanctuary and Protection Act 2008 establishes a sanctuary, which covers 8,234 square miles, taking in 16 islands and 145 islets and atolls, out to 12 miles offshore, specifically protecting its primarily reef manta ray population and its habitat. The Republic of Maldives bans exports of all ray species and their body parts and has created two marine protected areas (MPAs), specifically identified for protection because of their importance as areas of critical habitats for the Maldives population of reef manta rays and the occasional transient oceanic manta ray. In Western Australia, manta

rays (both species), whilst not targeted, are protected from any fishing (Fisheries Act) and disturbance or harassment (DEC Act) within marine parks only. Other range States protect Manta rays in relatively small marine park zones.

Effectiveness of these measures varies, with reports of illegal fishing of *Manta* spp. in Mexico and the Philippines (Graham et al. 2012, S. Heinrichs, pers. comm., Marshall et al. 2011c, GMA TV, May 2012). *Manta* spp. (primarily *M. alfredi*) are also targeted in the Komodo Marine Park, near Lamakera, Indonesia, despite regulations forbidding fishing (H. Dewar, pers. comm.).

#### 4.2 International protection status

At the last Conference of the Parties to the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) in March 2013, the proposal by the governments of Brazil, Colombia and Ecuador to add the genus *Manta*, including *M. birostris*, *M. alfredi* and any putative *Manta* species, to Appendix II was adopted. No other international protections are in place for *M. alfredi*.

*Manta alfredi* is listed as Vulnerable on the IUCN Red List of Threatened Species with a decreasing population trend.

#### 4.3 Additional protection status

At the 10th CMS CoP in November 2011 the proposal of Ecuador to add *M. birostris* to CMS Appendices I and II was adopted. While the recently separated species, *M. alfredi*, was not included in Ecuador's proposal, Norway, the host country, suggested that this species be listed at the next Conference of the Parties. Despite the Appendix I and II listing of *M. birostris*, *Manta* species have not yet been added to Annex I of the CMS Sharks MoU.

### 5. **Range states (See Annex I)**

### 6. **Comments from range states**

**Fiji:** Fiji does not have a targeted or subsistence fishery for Manta Ray species, but notes that the Manta ray species are caught as by-catch within the Purse seine Fisheries within the Western-Central Pacific Ocean. Manta Rays are largely not fished or harvested across the waters of the Fiji Islands, but used for ecotourism attractions in a number of targeted dive sites within Fiji's coastal reef and island systems. In Fiji, the local island systems that currently have Manta Ray dive tourism (primarily *M. alfredi*) are on the islands of Taveuni, Kadavu and the Lau groups. These rays migrate large distances across the Pacific and seem to come to Fiji's waters for abundant food & mating habitats.

### 7. **Additional remarks**

Countries across the South-west Pacific (include Tonga, Samoa, Vanuatu, Fiji, Cook Island, and others) have documented and observed how species of *Mobula*, *Manta* and other rays interact within their local coastal and associated areas of national jurisdictions, and clearly noted from dive operators in a number of the local island systems, that these species are one

of the big draw-cards for the dive and snorkel tourists to the region. Manta rays will receive protection under their CITES Appendix II listing in September-2014, but including them on the appendices of CMS would help support more comprehensive international protection. The devil ray populations within the South Pacific are also on the decline, and the rest of the South Pacific region would also be very supportive if Fiji were able to start some form of protection for these Manta and Mobula Rays. Even though the CMS is non-binding and voluntary, it is a strong indicator of countries showing willingness to take leadership in their conservation.

## 8. References

- Alava, E.R.Z., Dolumbaló, E.R., Yaptinchay, A.A., and Trono, R.B. 2002. Fishery and trade of whale sharks and manta rays in the Bohol Sea, Philippines. In: Fowler, S.L., Reed, T.M., Dipper, F.A. (eds) Elasmobranch Biodiversity, Conservation and Management: Proceedings of the International Seminar and Workshop. Sabah, Malaysia, July 1997, pp 132–148
- Amande, M.J., Ariz, J., Chassot, E., De Molina, A.D., Gaertner, D., Murua, H., Pianet, R., Ruiz, J., and Chavance, P. 2010. Bycatch of the European purse seine tuna fishery in the Atlantic Ocean for the 2003-2007 period. *Aquatic Living Resources*, 23(4): 353-362.
- Anderson, R.C., Adam, M.S., Kitchen-Wheeler, A., and Steven G. 2010. Extent and economic value of manta ray watching in the Maldives. *Tourism in Marine Environments*, 7(1): 15-27.
- Anderson, R.C., Adam, M.S., and Goes, J.I. 2011. From monsoons to mantas: seasonal distribution of *Manta alfredi* in the Maldives. *Fisheries Oceanography*, 20(2): 104-113.
- Barnes, R.H. 2005. Indigenous use and management of whales and other marine resources in East Flores and Lembata, Indonesia. *Senri Ethnological Studies*, 67: 77-85.
- Bigelow, H.B. and Schroeder, W.C. 1953. Sawfish, guitarfish, skates and rays. In: Bigelow, H.B. and Schroeder, W.C. (Eds) *Fishes of the Western North Atlantic, Part 2*. Sears Foundation for Marine Research, Yale University, New Haven, pp. 508-514.
- Booda, L. 1984. Manta ray wings, shark meat posing as scallops. *Sea Technology* 25(11): 71.
- Camhi, M.D., Valenti, S.V., Fordham, S.V., Fowler, S.L. and Gibson, C. 2009. The Conservation Status of Pelagic Sharks and Rays: Report of the IUCN Shark Specialist Group Pelagic Shark Red List Workshop. Newbury, UK: IUCN Species Survival Commission Shark Specialist Group, x +78 pp.
- Chavance, P., Amande, J.M., Pianet, R., Chassot, E., and Damiano, A. 2011. Bycatch and discards of the French Tuna Purse Seine Fishery during the 2003-2010 period estimated from observer data. IOTC-2011-WPEB07-23.
- Chin, A., Kyne, P.M. 2007. Vulnerability of chondrichthyan fishes of the Great Barrier Reef to climate change. In: *Climate Change and the Great Barrier Reef: A Vulnerability Assessment*, Johnson, J.E., and Marshall, P.A. (eds). Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Townsville, Australia. P 393-425.
- Clark, T.B. 2001. Population structure of *Manta birostris* (Chondrichthyes: Mobulidae) from the Pacific and Atlantic Oceans. MS thesis, Texas A&M University, Galveston, TX
- Clark, T.B. 2010. Abundance, Home Range, and Movement Patterns of Manta Rays, Doctoral thesis, University of Hawaii at Manoa
- Coan, A.L., Sakagawa, G.T., Prescott, D., Williams, P., Staish, K., and Yamasaki, G. 2000. The 1999 U.S. Central-Western Pacific Tropical Tuna Purse Seine Fishery. Document prepared for the annual meeting of parties to the South Pacific Regional Tuna Treaty 3-10 March 2000. LJ-00-10.
- Compagno, L.J.V. 1999. Checklist of living elasmobranchs. In: Hamlett, W.C. (ed). *Sharks, skates, and rays: the biology of elasmobranch fishes*. Maryland: John Hopkins University Press. p 471–498

- Compagno, L.J.V. and Last, P. 1999. Mobulidae. In: Capenter, K.E. and Niem, V.H. (eds), FAO species identification guide for fishery purposes. The living marine resources of the western Central Pacific (Volume 3. Batoid Fishes, Chimeras and Bony Fishes)
- Couturier, L.I.E., Dudgeon C.L., Pollock, K.H., Jaine, F.R.A., Bennett, M.B., Townsend, K.A., Weeks, S.J. and Richardson, A.J. 2014. Population Dynamics of the reef manta ray *Manta alfredi* in eastern Australia. Coral Reefs DOI 10.1007/s00338-014-1126-5
- Couturier, L.I.E., Marshall, A.D., Jaine, F.R.A., Kashiwagi, T., Pierce, S.J., Townsend, K.A., Weeks, S.J., Bennet, M.B., and Richardson, A.J. 2012. Biology, ecology and conservation of the Mobulidae. *Journal of Fish Biology*, 80: 1075-1119.
- Deakos, M.H. 2010. Ecology and social behavior of a resident manta ray (*Manta alfredi*) population off Maui, Hawai'i. PhD thesis, University of Hawai'i, Manoa, Hawai'i.
- Deakos, M., Baker, J., and Bejder, L. 2011. Characteristics of a manta ray (*Manta alfredi*) population off Maui, Hawaii, and implications for management. *Marine Ecology Progress Series*, 429: 245-260.
- Dewar, H. (2002). Preliminary report: Manta harvest in Lamakera. p. 3 p. Oceanside, USA: Report from the Pflieger Institute of Environmental Research and the Nature Conservancy.
- Dewar, H., Mous, P., Domeier, M., Muljadi, A., Pet, J., Whitty, J. 2008. Movements and site fidelity of the giant manta ray, *Manta birostris*, in the Komodo Marine Park, Indonesia. *Marine Biology*, Vol. 155, Number 2, 121-133.
- Donnelly, R., Neville, D., and Mous, P.J. 2003. Report on a rapid ecological assessment of the Raja Ampat Islands, Papua, Eastern Indonesia, held October 30 – November 22, 2002. The Nature Conservancy – Southeast Asia Center for Marine Protected Areas, 250 pp.
- Dulvy, N., Pardo, S., Simpfendorfer, C., Carlson, J. 2014. Diagnosing the dangerous demography of manta rays using life history theory. *PeerJ PrePrints* 162/v1: 1-26.
- Duinkerken, D.I. (2010) Movements and site fidelity of the reef manta ray, *Manta alfredi*, along the coast of southern Mozambique. MSc Utrecht University, The Netherlands.
- Essumang, D. 2010. First determination of the levels of platinum group metals in *Manta birostris* (Manta Ray) caught along the Ghanaian coastline. *Bulletin of Environmental Contamination and Toxicology*, 84(6): 720-725.
- Fernando, D. and Stevens, G. 2011 A study of Sri Lanka's manta and mobula ray fishery. The Manta Trust, 29 pp.
- Fowler, S. and S. Valenti/IUCN SSC Shark Specialist Group, 2007. Review of Migratory Chondrichthyan Fishes. CMS Technical Series No. 15.
- Germanov, E.S. and Marshall, A.D. (2014) Running the gauntlet: regional movement patterns of *Manta alfredi* through a complex of parks and fisheries. *PlosOne* In Press.
- GMA TV -- "Pangangatay ng manta ray at devil ray sa isla ng Pamilacan", Born to be Wild. Aired GMA TV Atlanta. 23 May 2012. Television.
- Graham, R.T., Witt, M.J., 2008. Site Fidelity and Movements of Juvenile Manta Rays in the Gulf of Mexico. AES Devil Ray Symposium, Joint Ichths and Herps Conference Presentation.
- Graham, R.T., Hickerson, E., Castellanos, D.W., Remolina, F., Maxwell, S. 2012. Satellite Tracking of Manta Rays Highlights Challenges to Their Conservation. *PLoS ONE* 7(5): e36834. Doi:10.1371/journal.pone.0036834
- Handwerk, B. 2010. Little-known Gulf manta ray affected by oil spill? *National Geographic News*, Published Oct. 15, 2010. <http://news.nationalgeographic.com/news/2010/10/101015-new-manta-ras-gulf-bp-oil-spill-science-animals/> accessed Sept. 1, 2011.
- Harding, M., and Beirwagen, S. 2009. Population research of *Manta birostris* in coastal waters surrounding Isla de la Plata, Ecuador.
- Heinrichs, S., O'Malley, M., Medd, H., and Hilton, P. 2011. Manta Ray of Hope: Global Threat to Manta and Mobula Rays. Manta Ray of Hope Project ([www.mantarayofhope.com](http://www.mantarayofhope.com)).


- Higgs, N.D., Gates, A.R., Jones, D.O.B. (2014) Fish food in the deep sea: revisiting the role of large fish falls. *Plos One* 9(5):e96016.
- Hilton, P. 2011. East Asia Market Investigation. Manta Ray of Hope, 49pp.
- Homma, K., Maruyama, T., Itoh, T., Ishihara, H., and Uchida, S. 1999. Biology of the manta ray, *Manta birostris* Walbaum, in the Indo-Pacific. In: Seret, B. and Sire, J.Y. (eds) Indo-Pacific fish biology: Proc 5th Int Conf Indo-Pacific Fishes, Noumea, 1997. Ichthyological Society of France, Paris, p 209–216
- Jaine FRA, Couturier LIE, Weeks SJ, Townsend KA, Bennett MB, et al. (2012) When Giants Turn Up: Sighting Trends, Environmental Influences and Habitat Use of the Manta Ray *Manta alfredi* at a Coral Reef. *PLoS One* 7: e46170.
- Jaine FRA, Rohner CA, Weeks SJ, Couturier LIE, Bennett MB, et al. (2014) Movements and habitat use of reef manta rays off eastern Australia: Offshore excursions, deep diving and eddy affinity revealed by satellite telemetry. *Mar Ecol Prog DOI: 10.3354/meps10910*.
- Kashiwagi, T. Marshall, A. D., Bennett, M. B., and Ovenden, J. R. 2011. Habitat segregation and mosaic sympatry of the two species of manta ray in the Indian and Pacific Oceans: *Manta alfredi* and *M. birostris*. *Marine Biodiversity Records*: 1-8.
- Kashiwagi, T., Marshall, A. D., Bennett, M.B., and Ovenden, J.R. 2012. The genetic signature of recent speciation in manta rays (*Manta alfredi* and *M. birostris*). *Molecular Phylogenetics and Evolution*, 64(1): 212-218.
- Kitchen-Wheeler, A. 2008. Migration behaviour of the Giant Manta (*Manta birostris*) in the Central Maldives Atolls. Paper presented at the 2008 Joint Meeting of Ichthyologists and herpetologists, Montreal, Canada.
- Kitchen-Wheeler, A. 2010. Visual identification of individual manta ray (*Manta alfredi*) in the Maldives Islands, Western Indian Ocean. *Marine Biology Research*, 6(4):351-363
- KMP (Komodo Manta Project). 2011. Manta population estimations from photographs. Unpublished Data.
- Lack, M and Sant, G. 2009. Trends in global shark catch and recent developments in management. *TRAFFIC International*, 33 pp.
- Marshall, A.D., Pierce, S.J., Bennett, M.B., 2008. Morphological measurements of manta rays (*Manta birostris*) with a description of a foetus from the east coast of Southern Africa. *Zootaxa*, 1717: 24-30.
- Marshall, A. D. 2009. Biology and population ecology of *Manta birostris* in southern Mozambique. PhD Thesis, University of Queensland
- Marshall, A., Compagno, L., Bennett, M. 2009. Redescription of the genus *Manta* with resurrection of *Manta alfredi* (Krefft, 1868) (Chondrichthyes; Myliobatoidei; Mobulidae). *Zootaxa* 2301: 1-28.
- Marshall, A.D., Holmer, J., Brunnschweiler, J.M. and Pierce, S.J. 2010. Size structure and migratory behaviour of a photographically identified population of *Manta birostris* in southern Mozambique.
- Marshall, A.D., Dudgeon, C.L. and Bennett, M.B. 2011a. Size and structure of a photographically identified population of manta rays *Manta alfredi* in southern Mozambique. *Marine Biology*, 158 (5): 1111-1124.
- Marshall, A., Kashiwagi, T., Bennett, M.B., Deakos, M., Stevens, G., McGregor, F., Clark, T., Ishihara, H. & Sato, K. 2011b. *Manta alfredi*. In: IUCN 2011. IUCN Red List of Threatened Species. Version 2011.2. <[www.iucnredlist.org](http://www.iucnredlist.org)>.
- Marshall, A., Bennett, M.B., Kodja, G., Hinojosa-Alvarez, S., Galvan-Magana, F., Harding, M., Stevens, G. & Kashiwagi, T. 2011c. *Manta birostris*. In: IUCN 2011. IUCN Red List of Threatened Species. Version 2011.2. <[www.iucnredlist.org](http://www.iucnredlist.org)>.
- McCauley, D.J., DeSalles, P.A., Young, H.S., Dunbar, R.B., Dirzo, R., Mills, M.M., and Micheli, F. 2012. From wing to wing: the persistence of long ecological interaction chains in less-

- disturbed ecosystems. *Scientific Reports*, 2: 409.
- McGregor, F. 2009. The Manta Rays of Ningaloo Reef: baseline population and foraging ecology. Presentation, Murdoch University.
- Mohanraj, G., Rajapackiam, S., Mohan, S., Batcha, H., and Gomathy, S. 2009. Status of elasmobranchs fishery in Chennai, India. *Asian Fisheries Science*, 22: 607-615.
- Molony, B. 2005. Estimates of the mortality of non-target species with an initial focus on seabirds, turtles and sharks. 1<sup>st</sup> Meeting of the Scientific Committee of the Western and Central Pacific Fisheries Commission, 84 pp.
- MPRF (Manta Pacific Research Foundation). 2011. Manta ray photo-identification catalogue. [www.mantapacific.org/identification/index.html](http://www.mantapacific.org/identification/index.html). Accessed September 14, 2011.
- Notarbartolo di Sciara, G. and Hillyer, E.V. 1989. Mobulid rays off eastern Venezuela (Chnodrichthyes, Mobulidae). *Copeia*, 3: 607-614.
- Notarbartolo di Sciara, G. 1995. What future for manta rays? *Shark News*, 5: 1.
- Notarbartolo di Sciara, G. 2005. Giant devilray or devil ray *Mobula mobular* (Bonnaterre, 1788). In: *Sharks, Rays and Chimaeras: The Status of Chaondrichthyan Fishes*. Fowler, S.L., Cavanagh, R.D., Camhi, M., Burgess, G.H., Caillet, G.M., Fordham, S.V., Simpendorfer, C.A., and Musick, J.A. (eds.). Gland, Switzerland and Cambridge, UK: IUCN/SSC Shark Specialist Group, pp. 356-357.
- O'Malley M.P., Lee-Brooks K., Medd H.B. 2013. The Global Economic Impact of Manta Ray Watching Tourism. *PLoS ONE* 8(5): e65051.
- Papastamatiou, Y., DeSalles, P., & McCauley, D., 2012. Area-restricted searching by manta rays and their response to spatial scale in lagoon habitats. *Marine Ecology Progress Series*, 456, 233-244. doi:10.3354/meps09721
- Paulin, C.D., Habib, G., Carey, C.L., Swanson, P.M., Voss, G.J. 1982. New records of *Mobula japonica* and *Masturus lanceolatus*, and further records of *Luvaris imperialis* (Pisces: Mobulidae, Molidae, Louvaridae) from New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 16: 11-17.
- Perez, J.A.A. and Wahrlich, R. 2005. A bycatch assessment of the gillnet monkfish *Lophius gastrophysus* fishery off southern Brazil. *Fisheries Research*, 72: 81-95.
- Pianet, R., Chavance, P., Murua, H., Delgado de Molina, A. 2010. Quantitative estimates of the by-catches of the main species of the purse seine fleet in the Indian Ocean, 2003-2008. Indian Ocean Tuna Commission, WPEB-21.
- Pillai, S.K. 1998. A note on giant devil ray *Mobula diabolus* caught in Vizhinjam. *Marine Fisheries Information Service, Technical and Extension Series*, 152: 14-15.
- Planeta Oceano 2011. Preliminary report of the state of coastal mobulid fisheries in Peru.
- Poortvliet, M., Galvan-Magana, F., Bernardi, G., Croll, D.A., and Olsen, J.L. 2011. Isolation and characterization of twelve microsatellite loci for the Japanese Devilray (*Mobula japonica*). *Conservation Genetics Resource*. 3: 733-735.
- Rajapackiam, S. Mohan, S. and Rudramurthy, N. 2007. Utilization of gill rakers of lesser devil ray *Mobula diabolus* – a new fish byproduct. *Marine Fisheries Information Service, Technical and Extension Series*, 191: 22-23.
- Raje, S. G., Sivakami, S., Mohanraj, G., Manojkumar, P.P., Raju, A. and Joshi, K.K. 2007. An atlas on the Elasmobranch fishery resources of India. CMFRI Special Publication, 95. pp. 1-253.
- Rohner, C., Pierce, S., Marshall, A., Weeks, S., Bennett, M., Richardson, A. 2013. Trends in sightings and environmental influences on a coastal aggregation of manta rays and whale sharks. *Marine Ecology Progress Series*, 482: 153-168.
- Romanov, E.V. 2002. Bycatch in the tuna purse-seine fisheries of the western Indian Ocean. *Fishery Bulletin*, 100(1): 90-105
- Rubin, R.D. 2002. Manta Rays: not all black and white, *Shark Focus*, 15: 4-5.

- Stevens, G., 2011, Field Guide to the Identification of Mobulid Rays (Mobulidae): Indo-West Pacific. The Manta Trust. 19 pp.
- Thomas, P., 1994, Preying on Mantas: After Divers Videotape Slaughter, Officials Enact Regulation to Aid Rays off Mexican Island., Los Angeles Times, 13 April.
- Tomita, T., Toda, M., Ueda, K., Uchida, S., Nakaya, K. 2012. Live-bearing manta ray: how the embryo acquires oxygen without placenta and umbilical cord. *Biol. Lett.* Published online 6 June 2012, doi: 10.1098/rsbl.2012.0288.
- Uchida, S. 1994. Manta Ray, basic data for the Japanese threatened wild water organisms (pp.152-159). Tokyo, Japan: Fishery Agency of Japan.
- Whitcraft, S., O'Malley, M., Hilton, P. 2014. The Continuing Threat to Manta and Mobula Rays: 2013-14 Market Surveys, Guangzhou, China. WildAid, San Francisco, CA
- White, W.T., Clark, T.B., Smith, W.D. & Bizzarro, J.J. 2006. *Mobula japonica*. In: IUCN 2011. IUCN Red List of Threatened Species. Version 2011.2. <[www.iucnredlist.org](http://www.iucnredlist.org)>
- White, W.T., Clark, T.B., Smith, W.D. & Bizzarro, J.J. 2006. *Mobula japonica*. In: IUCN 2011. IUCN Red List of Threatened Species. Version 2011.2. <[www.iucnredlist.org](http://www.iucnredlist.org)>
- White, W. T., Giles, J., Dharmadi, and Potter, I. C. 2006. Data on the bycatch fishery and reproductive biology of mobulid rays (Myliobatiformes) in Indonesia. *Fisheries Research*, 82(1-3), 65-73.
- White, W., and Kyne, P. 2010. The status of chondrichthyan conservation in the Indo-Australasian region. *Journal of Fish Biology*, 76(9), 2090-2117
- Young, N. 2001. An analysis of the trends in by-catch of turtle species, angelsharks and batoid species protective gillnets off KwaZulu-Natal, South Africa. Msc. Thesis, University of Reading.
- Zeeberg, J., Corten, A., and de Graaf, E. 2006. Bycatch and release of pelagic megafauna in industrial trawler fisheries off Northwest Africa. *Fisheries Research*, 78: 186-195.

ANNEX I. Distribution Map and Table

**Worldwide Distribution of *Manta alfredi***



**Distribution Table – *Manta alfredi* Range States and FAO Fisheries Areas**

<b>Range States and FAO Fisheries Areas</b>	<b><i>Manta alfredi</i></b>		
FAO Fisheries Areas	51, 57, 71, 77, 81		
South Africa (Eastern Cape Province, KwaZulu-Natal, Western Cape Province)	x		
Mozambique	x		
Madagascar (Nosy Be)	x		
Comoros - Mayotte (France)	x		
Egypt - Sinai (African part)	x		
Saudi Arabia	x		
Sudan	x		
Djibouti	x		
Yemen	x		
Oman	x		
Seychelles (Inner Island Group, Amirante Islands, Farquhar Group, Aldabra Group)	x		
Chagos Archipelago (British Indian Ocean Territory)	x		
Maldives	x		
Pakistan	x		
India (Lakshadweep & Andaman Is. - <i>M. alfredi</i> & <i>M. birostris</i> - Andhra Pradesh, Goa, Gujarat, Kerala, Maharashtra, Tamil Nadu - <i>M. birostris</i> only)	x		
Thailand	x		
Malaysia	x		
Indonesia (Sumatra, Bali, Komodo, Flores, Irian Jaya, Java, Lombok, Alor, Borneo, Sulawesi)	x		
		Cocos (Keeling) Islands (Australia)	x
		Australia (New South Wales, Northern Territory, Queensland, Western Australia)	x
		Philippines (Monad Shoal, Tubbataha Reef, Pamilacan, Apo Reef, Gigdup Shoal, Ticau & Masbate)	x
		Ryukyu & Nampo-shoto Archipelagos' (Japan)	x
		Northern Mariana Islands (Saipan) & Guam (US)	x
		Federated States of Micronesia (Yap, Pohnpei)	x
		Palau	x
		Papua New Guinea (Bismarck Archipelago, North Solomon's, Main Island Group)	x
		Solomon Islands	x
		New Caledonia (France)	x
		Vanuatu	x
		Marshall Islands	x
		Fiji	x
		Tuvalu	x
		Tonga	x
		Cook Islands	x
		Kiribati (Christmas Island)	x
		Line Islands - Jarvis, Palmyra & Kingman (US)	x
		Hawaiian Islands (US)	x
		French Polynesia - Society, Marquises & Tuamotu Islands	x

**ANNEX II. *Manta alfredi* legal protective measures**

<b><i>Manta alfredi</i> legal protective measures</b>		
<b>Location</b>	<b>Species</b>	<b>Legal Protection / Conservation Measure</b>
<b>International</b>		
CITES Appendix II	<i>Manta spp.</i>	Listing of the genus <i>Manta</i> on Appendix II of the Convention on International Trade in Endangered Species (CITES), 2013
<b>National</b>		
Indonesia	<i>Manta spp.</i>	KepMen National Protective Legislation, 2014
Maldives	<i>Manta spp.</i>	Exports of all ray products banned 1995
Yap (FSM)	<i>Manta spp.</i>	Manta Ray Sanctuary and Protection Act 2008
<b>State</b>		
Florida, USA	Genus <i>Manta</i>	FL Admin Code 68B-44.008 – no harvest
Guam, USA Territory	All ray species	Bill 44-31 prohibiting possession/sale/trade in ray parts 2011
Hawaii, USA	<i>Manta spp.</i>	H.B. 366 2009 – no harvest or trade
Raja Ampat Regency, Indonesia	<i>Manta spp.</i>	Shark and Ray Sanctuary Bupati Decree 2010

## **ADDITIONAL INFORMATION PROVIDED BY THE GOVERNMENT OF FIJI TO THE PROPOSAL FOR THE INCLUSION OF MANTA ALFREDI IN APPENDIX I AND II**

During the meeting of the CMS Scientific Council in July, members requested that Fiji provide additional information on the benefits of listing in the Appendices of the Convention to this species.

Appendix I listing would encourage Range State Parties where *M. alfredi* are targeted to seek help with implementing measures to protect this species and enable artisanal fishers to benefit from the much more lucrative and sustainable tourism revenues this species generates. For example in Mozambique, *Manta alfredi* and the closely related *Manta birostris* (already included on Appendix I) contribute over US\$13 million annually to tourism revenues. However, off Praia do Tofo, an important manta and Whale shark tourism area in southern Mozambique, artisanal fishers opportunistically target mantas for their low value meat. In this area, Rohner et al. (2013) observed a decline in *Manta alfredi* abundance of 88 per cent over only eight years (less than a third of one generation period for this species).

In addition, new evidence of growing demand for manta and mobula gill plates and expansion of this trade threatens to motivate the emergence of new, targeted manta fisheries in Range States where *M. alfredi* are currently not protected. CMS Parties that are Range States of *M. alfredi*, which currently do not have protection measures in place for *M. alfredi*, include Australia, Comoros, Cook Islands, Djibouti, Egypt, Fiji, France (Mayotte and New Caledonia), India, Madagascar, Mozambique, Pakistan, Palau, Philippines, Saudi Arabia, Seychelles, South Africa and Yemen.

The CMS Appendix I listing, therefore, will help this species by curbing existing targeted fisheries and preventing emergence of new fisheries in response to demand from the mobulid gill plate trade.