



**Convention on Migratory Species
Memorandum of Understanding for the
Conservation of Cetaceans and their
Habitats in the Pacific Islands Region**

Distr: General

UNEP/CMS/PIC2/Inf.6-04
15 July 2009

Agenda Item 6.3

SECOND MEETING OF THE SIGNATORIES
Auckland, New Zealand, 28-29 July 2009

**IUCN PROPOSAL
FOR SEPARATE LISTING FOR OCEANIA SUB-POPULATION OF HUMPBACKS
(*Megaptera novaeangliae*)**

'Submission by South Pacific Whale Research Consortium et al to IUCN (November 2008) on the classification of humpback whales, which resulted in the Oceania population of humpback whales being reclassified from Vulnerable to Endangered'

Megaptera novaeangliae (Borowski, 1781) (Oceania sub-population)

Humpback whale (Oceania sub-population)

Taxonomic Notes

The humpback whale is a cosmopolitan species and is regarded as monotypic (Rice 1998; Clapham & Mead 1999). No sub-species are recognised. Phylogenetic analyses reveal that populations in different ocean basins do not represent distinct clades, but some maternal lineages can be traced back across hemispheric boundaries, indicating some level of inter-hemispheric gene flow, perhaps in the distant past (Baker *et al.* 1994).

Population structure

The International Whaling Commission (IWC) currently recognises four breeding stocks around the South Pacific and Australia based on Discovery mark recoveries, demographic isolation, and genetic differentiation (Olavarria *et al.* 2007):

- 1 north of feeding area IV (referred to as Stock D including Western Australia),
- 1 north of feeding Area V (referred to as Stock E including Eastern Australia, New Caledonia and Tonga),
- 1 north of feeding Area VI (Stock F including Cook Islands and French Polynesia), and
- 1 north of feeding Area I (Stock G including Colombia).

The IWC also recognises further stock sub-division of breeding stock E and F into sub-stocks supported by demographic isolation and genetic differentiation (Olavarria *et al.* 2006). Breeding stock E is sub-divided into E1 (Eastern Australia), E2 (New Caledonia) and E3 (Tonga) and breeding stock F into F1 (Cook Islands) and F2 (French Polynesia). These breeding stocks are shown in Appendix 1. We use the terminology breeding stock (e.g. D, E, F, G) to refer to breeding stocks based on demographic isolation and genetic differentiation, and sub-stock to refer to sub-divisions within these breeding stocks, as currently recognised by the IWC (e.g. E1, E2, E3, F1, F2).

Olavarria *et al.* (2007) found significant differentiation of maternally inherited mitochondrial (mt) DNA at both the haplotype and nucleotide level ($F_{ST} = 0.033$; $\Phi_{ST} = 0.022$), between breeding stocks D, G and four of the Oceania sub-stocks (E2, E3, F1, F2). When sub-stock E1 is included in this comparison (Olavarria *et al.* (2006)), the estimated differentiation among stocks by F_{ST} is ~ 0.02 . Based on standard population genetic models (e.g. Wright 1978, Waples & Gaggiotti 2006), F_{ST} values of 0.01 correspond to approximately 25 migrant individuals per generation (or less than one migrant individual per year in the case of humpback whales and other long lived mammals). These breeding stock boundaries, and the sub-stocks within them, are also supported by the analysis of movements by individuals based on photo-identification and microsatellite genotyping (DNA profiling).

An extensive comparison of photo-identification catalogues by sub-stock, found only four matches between the migratory corridor of East Australia (E1 represented by Hervey Bay and Byron Bay, with a catalogue size of 1,242 individuals) and breeding grounds in Oceania (E2, E3, F1 and F2, with a catalogue size of 679 individuals) (Garrigue *et al.* 2007). This level of interchange is surprisingly small, given the relatively large catalogues used in the comparison, and provides strong evidence for sub-division within Breeding Stock E (Garrigue *et al.* 2007).

An additional photo-identification comparison among regions of Oceania (E2, E3, F1 and F2 with a combined catalogue size of 679) documented 20 records of interchange, mostly between neighbouring regions (Garrigue *et al.* 2006). Overall, the limited movement of individuals between adjacent sites within Oceania is consistent with the significant (but low) level of differentiation observed in mtDNA from these regions (Olavarria *et al.* 2007) and suggests that humpback whales wintering in E2, E3, F1 and F2 are demographically independent and should be recognised as individual management stocks (Garrigue *et al.* 2006).

Comparisons of historical sighting data and whaling records (Dawbin 1956, 1959, 1964) with recent sighting survey data from New Zealand, Fiji and Norfolk Island demonstrate a lack of (or at the very least a slow) recovery at these sites (Childerhouse & Gibbs 2006; Gibbs *et al.* 2006; Paton *et al.* 2006; Oosterman & Whicker 2008). These surveys returned to the same look out sites used by Dr. W. Dawbin and replicated his earlier surveys as closely as possible. Results from these re-surveys include (i) sighting rates in Fiji over the period 1956-58 were between 0.15-0.58 whales per hour and were significantly higher than equivalent sighting rates observed of between 0.01-0.03 in 2002-03 (Paton *et al.* 2006) and, (ii) surveys in New Zealand indicate that between 2004-2006 sightings were 29% of what there were in 1960 (Childerhouse & Gibbs 2006). It is important to note that the baseline data for these surveys in the 1950s and 1960s were from populations that had already been whaled for more than 50 years. It is not possible to directly assess the rates of increase for these sites but what is clear is that any population increases appear to be lacking or very low. In contrast, the East Australian stock is increasing at 10-11% per annum (Noad *et al.* 2006).

These indications of demographic independence are likely sufficiently strong to provide evidence for further sub-populations within the Oceania, however, such partitioning presents difficulties in assessing population status (discussed below) that have not been overcome at present. Furthermore, problems with the allocation of commercial catches on the feeding grounds to the appropriate sub-stock breeding area make the assessment even more challenging. Given it is not possible to assess the status of each sub-stock, we have therefore used a model that can assess the South Pacific as though it is a single stock (i.e. E and F).

In conclusion, the presently recognised IWC stock and sub-stock boundaries are consistent with available evidence. With respect to the South Pacific, the relevant sub-stock divisions are East Australia (E1), New Caledonia (E2), Tonga (E3), Cook Islands

(F1), French Polynesia (F2), and Colombia (G). The taxon assessed here is, therefore, called the Oceania sub-population, which consists of IWC breeding stocks E and F as a distinct sub-population of humpback whales. It should be identified separately based on population isolation and a demonstrated high level of depletion (see below). We propose this sub-population specifically for the purposes of the IUCN threat ranking process as it is consistent with the existing IWC recognised breeding stock boundaries.

Geographic Range Information

Humpback whales have a global distribution. Individual humpbacks have been observed to travel more than 8000km between their high-latitude summer feeding grounds and winter mating and calving range in tropical waters (Rasmussen *et al.* 2007). The Oceania sub-population is delineated by its breeding range, with approximate boundaries in the west at 160°E (between Australia and New Caledonia), in the east at 120°W (between French Polynesia and South America), in the north at the equator at 0°S, and in the south to approximately 30°S.

During the austral autumn and winter, humpback whales in Oceania are spread across lower latitudes from approximately 30°S northwards to the equator. The South Pacific is a vast area with thousands of islands and there has not yet been a comprehensive survey of the entire region. However, localised research by members of the South Pacific Whale Research Consortium (SPWRC 2008)) has identified many island groups whose waters are host to humpback whales. During austral spring and summer, humpbacks travel to Antarctic feeding grounds. These linkages have been demonstrated through Discovery tagging, photo-identification and, most recently, genotype matching and satellite telemetry (Mackintosh 1942; Chittleborough 1965; Dawbin 1966; Mikhalev 2000; Franklin *et al.* 2007).

Range Countries and territories

Country and Territory Names: American Samoa (United States of America), Australia, Cook Islands, Federated States of Micronesia, Fiji, French Polynesia (France), Guam (United States of America), Republic of Kiribati, Marshall Islands, Nauru, New Caledonia (France), New Zealand, Niue, Northern Mariana Islands, Palau, Papua New Guinea, Samoa, Solomon Islands, Tokelau, Kingdom of Tonga, Tuvalu, Vanuatu, Wallis and Futuna

Aquatic Regions: Pacific-South

Population Information

The following population estimates are available:

(i) SPWRC (2006) provided a preliminary mark-recapture estimate from photo-identification of the combined population size for E2 (New Caledonia), E3 (Tonga) and F (French Polynesia) of 3,827 (CV = 0.12) for the period 1999-2004. There are no

estimates of rate of increase available for this area but it was noted that there was little indication of trend in abundance over the survey period (SPWRC 2006).

(ii) Noad *et al.* (2006) estimated from land-based sighting surveys that population size of E1 (Eastern Australia) was 7,090 (95% CI \pm 660) for 2004 with an annual rate of increase of 10.6 (95% CI \pm 0.5%) for 1987 – 2004.

The IWC is presently engaged in a Comprehensive Assessment of Southern Hemisphere humpback whales and research on the South Pacific breeding stocks of E1, E2, E3, and F are ongoing. The IWC (2007) Comprehensive Assessment of Southern Hemisphere Humpback workshop in 2006 agreed that, “*the situation for Breeding Stocks E and F is complex and currently unresolved, and therefore that it was not possible to construct stock structure hypotheses for assessment modelling, particularly with respect to the assignment to Breeding Stocks of catches taken on the feeding grounds*”.

For example, while east Australia and New Caledonia (E1 and E2) are within the longitudinal boundaries of Antarctic Area V, and French Polynesia and the Cook Islands (F) are within the longitudinal boundaries of Area VI, Tonga (E3) falls close to the boundary between the two Areas. Thus, in the current assessment, the approach of pooling demographically independent sub-populations was necessary for practical reasons to develop catch allocation scenarios. However, this approach is likely to be conservative in ignoring potential differences in variable rates of recovery from the regional impacts of whaling. Soviet whaling on the Antarctic feeding grounds in the early sixties was extremely intense, with over 27,300 whales taken during two summers (1959-1961) alone. Maternal site fidelity together with a hunt concentrated both in time and space may have resulted in more extreme declines in some of the far-flung wintering stocks of the Southwestern Pacific.

Jackson *et al.* (2006) explored a number of catch allocation scenarios for the combined sub-stocks of Oceania and east Australia. In their combined assessment of sub-stocks E1, E2, E3 and F, median population recovery toward historical levels in 2005 was estimated at between 15.9-24.8% (95% probability intervals (PI) 11.1-30.5%; prior population growth rate mean = 6.7% after Branch *et al.* (2004)). The most appropriate interpolation between these two recovery estimates depended on the degree of interchange between east Australia and Oceania (15.9% is complete interchange, 24.8% is no interchange). Recent photo-identification surveys (Garrigue *et al.* 2007) indicate that interchange between these regions is relatively low, suggesting that the ‘no interchange’ scenario may be more appropriate for the region. Under this interchange scenario, estimated abundance in 1942 was 41,356 (95% PI 36,800-53,580). Recovery of the population three generations later (in 2005) is 26.6% (95% PI 18.2-33.5%) relative to 1942. This is using an estimate of 21.5 years/generation (Taylor *et al.* 2007).

Habitat and Ecology Information

Humpback whales have been recorded across most of the South Pacific, although densities vary from large numbers in East Australia to very low numbers in Fiji (in E3) and parts of French Polynesia. They are regularly found around island groups but are also

in open water away from islands. Humpbacks have been recorded throughout the southern ocean including south to the ice edge and in the Ross Sea.

Little is known regarding life history parameters for the Oceania population of humpback whales, although it is assumed that these rates are similar to those described from whaling records in Australia and New Zealand (Dawbin 1956, 1964, 1966; Chittleborough 1965). One rate that has been preliminarily investigated in the region is calving interval, which is approximately 2-3 years (consistent with that reported from other oceans). The diet of these humpback whales consists mainly of krill, which they consume while in Antarctic waters. They are not known to feed while in tropical breeding grounds.

Threat Information

During the last 2 centuries, humpback whales have been hunted intensively, especially in the Southern Hemisphere, where it was estimated that populations were reduced to a few percent of their pre-exploitation abundance (Chapman 1974). Based on catch records corrected for illegal Soviet whaling, a total of more than 200 000 humpback whales were killed in the Southern Hemisphere from 1904 to 1980 (Clapham & Baker 2002). Catches during the 19th century in the South Pacific by American whalers were made mainly during winter months in 3 tropical breeding grounds: off Colombia and Ecuador, around the Tongan archipelago, and northwest of New Caledonia (Townsend 1935; Mackintosh 1942). During the 20th century, humpback whales were hunted along their migratory corridors, such as along the coasts of New Zealand and Australia, and more intensively in their feeding areas in sub-Antarctic and Antarctic waters (Mackintosh 1942, 1965). The IWC gave legal protection to humpback whales from commercial whaling in 1966 but they continued to be killed illegally by whalers from the Soviet Union until 1972. Illegal Soviet takes of 25,000 humpback whales in two seasons (1959/60 and 1960/61) precipitated a population crash and the closure of land stations in Australia and New Zealand (Mikhalev 2000; Clapham *et al.* 2005).

Recently, Japan proposed to kill 50 humpback whales as part of its programme of scientific research under special permit (scientific whaling) in the IWC management areas IV and V in the Antarctic. Areas IV and V have demonstrated links with breeding stocks E. Japan postponed its proposed catch in the 2007/08 season but have not removed humpback whales from its future whaling programme. The continuation of this programme has the potential to slow the recovery of the Oceania sub-population.

Mortality of humpback whales due to entanglements in fishing gear and collisions with ship have been reported in the Southern Hemisphere (IWC 2001). Entanglement of humpback whales in pot lines occurs in both New Zealand and Australia. There is little information from around the rest of the South Pacific but a humpback mother (with calf) was reported entangled in a longline in 2007 (N. Hauser, reported in SPWRC 2008) and another humpback was struck and killed by a vessel in 1999 in Tonga (Diver 2004).

Conservation Measures

Although humpback whales have been legally protected from commercial whaling since 1966, they can still be killed for the purposes of scientific research under Article VIII of the International Convention for the Regulation of Whaling. The IWC's Southern Ocean Whale Sanctuary (e.g. the northern boundary of this Sanctuary follows the 40°S parallel of latitude except in the Indian Ocean sector where it joins the southern boundary of that sanctuary at 55°S, and around South America and into the South Pacific where the boundary is at 60°S) provides an additional layer of protection to humpback whales while on their summer feeding grounds in Antarctica although whales inside the Sanctuary can still be killed under Article VIII.

At present, more than 12 million km² of EEZs of more than a dozen South Pacific countries and territories have been designated as whale sanctuaries. This provides protection from commercial whaling for humpback whales in some of their breeding areas.

New Zealand and Australia have active disentanglement programmes to release any humpback whales captured in fishing gear.

Red List Assessment

Assessment: Endangered A1 ad

Rationale: The Oceania sub-population (as characterised by IWC breeding stocks E and F) is genetically and demographically isolated from their adjacent breeding stocks D (Western Australia) and G (Colombia). The final assessment of the Oceania sub-population has demonstrated that it is likely to have declined >70% in the last 3 generations (e.g. since 1942 using 21.5 years/generation (Taylor *et al.* 2007)). The estimated level of decline is based on the Jackson *et al.* (2006) combined assessment of E and F in which median population recovery towards historic levels in 2005 was estimated at 26.6% (95% probability intervals (PI) 18.2-33.5%). A 26.6% recovery equates to a 73.4% decline from estimated abundance prior to whaling. The Guidelines state, "In general, when uncertainty leads to wide variation in the results of assessments, the range of possible outcomes should be specified. A single category must be chosen and the basis for the decision should be documented; it should be both precautionary and credible". The range of possible outcomes spans both the Endangered and Vulnerable categories, however, the median outcome, which is consistent with guidelines needing to be both precautionary and credible, place this population in the Endangered category. It is likely that with additional analyses, the Oceania sub-population will be split into additional sub-populations that are experiencing different levels of risk, but until historical kills can be adequately addressed, assessing E and F stocks together represents the best available science.

Previous assessment: None.

Assessors: Simon Childerhouse, Jennifer Jackson, Scott Baker, Nick Gales, Philip Clapham, Robert Brownell

Evaluators: Randy Reeves, Steve Reilly, Howard Rosenbaum, Barbara Taylor

References

- Baker C.S., Slade, R. W., Bannister, J. L., Abernethy, R. B., Weinrich, M. T., Lien, J., Urbán, J., Corkeron, P., Calambokidis, J., Vasquez, O. and Palumbi, S. R. 1994. Hierarchical structure of mitochondrial DNA gene flow among humpback whales *Megaptera novaeangliae*, world-wide. *Mol. Ecol.* (4):313-271
- Baker, C.S.; Garrigue, C.; Constantine, R.; Madon, B.; Poole, M.; Hauser, N.; Clapham, P.; Donoghue, M.; Russell, K.; Paton, D.; Mattila, D. 2006. Abundance of humpback whales in Oceania (South Pacific), 1999 to 2004. Paper SC/A06/HW51 presented to IWC Comprehensive Assessment of Southern Hemisphere Humpback Whales Workshop 2006, Hobart, Australia.
- Branch T. A., K. Matsuoka, and T. Miyashita. 2004. Evidence for increases in Antarctic blue whales based on Bayesian modelling. *Marine Mammal Science* 20:726-754.
- Chapman, D.G. 1974. Status of Antarctic rorqual stocks. Pages 218-238 in W.E. Schevill, (ed.) *The whale problem*. Harvard University Press, Cambridge.
- Childerhouse, S.; Gibbs, N. 2006. Preliminary Report for the Cook Strait Humpback Whale Survey 2006. Unpublished Report to the Department of Conservation, New Zealand. 6 p.
- Chittleborough, R.G. 1965. Dynamics of two populations of humpback whales, *Megaptera novaeangliae* (Borowski). *Aust. J. Mar. and Freshw. Res.* 16:33-128.
- Clapham, P.J.; Baker, C.S. 2002. Modern whaling. Pages 1328-1332 in W.F. Perrin, B. Würsig and J.G.M. Thewissen, (eds.) *Encyclopedia of Marine Mammals*. Academic Press, New York.
- Clapham, P.J.; Aguilar, A.; Hatch, L. 2008. Determining spatial and temporal scales for management: lessons from whaling. *Marine Mammal Science* 24: 183-201.
- Clapham, P., Mikhalev, Yu., Franklin, W., Paton, D., Baker, C.S. & Brownell, R.L. Jr. 2005. Catches of humpback whales in the Southern Ocean, 1947-1973. Paper SC/57/SH6 presented to the International Whaling Commission Scientific Committee.
- Clapham P.J. and Mead J.G. 1999. *Megaptera novaeangliae*. *Mammalian Species* 604: 1-9.
- Dawbin, W.H.; Falla, R.A. 1949. A contribution to the study of the humpback whale based on observations at New Zealand shore stations. Pages 373-382 7th Pacific Science Congress.
- Dawbin, W.H. 1954. Maori Whaling. *The Norwegian Whaling Gazette* 8:433-445.
- Dawbin, W.H. 1956. Whale marking in South Pacific waters. *Norsk Hvalfangsttid* 45:485-508.
- Dawbin, W.H. 1956. The migration of humpback whales as they pass the New Zealand Coast. *Transactions of the Royal Society of New Zealand* 84:147-196.
- Dawbin, W.H. 1959. New Zealand and South Pacific whale marking and recoveries to the end of 1958. *Norsk Hvalfangsttid* 48:213-238.
- Dawbin, W.H. 1964. Movements of humpback whales marked in the southwest Pacific Ocean 1952 to 1962. *Norsk Hvalfangsttid* 53:68-78.
- Dawbin, W.H. 1966. The seasonal migratory cycle of humpback whales. Pages 145-171 in K.S. Norris, ed. *Whales, dolphins and porpoises*. University of California Press, Berkeley.
- Diver Co. Ltd. 2004. *Swimming with Giants*. Diver Magazine. UK
- Franklin, T.; Smith, F.; Gibbs, N.; Childerhouse, S.; Burns, D.; Paton, D.; Franklin, W.; Baker, C.S.; Clapham, P. 2007. Migratory movements of humpback whales (*Megaptera novaeangliae*) between eastern Australia and the Balleny Islands, Antarctica, confirmed by photo-identification. Paper SC/59/SH18 presented to the Scientific Committee of the International Whaling Commission Annual Meeting 2007, Anchorage, Alaska.
- Garrigue, C.; Baker, C.S.; Constantine, R.; Poole, M.; Hauser, N.; Clapham, P.; Donoghue, M.; Russell, K.; Paton, D.; Mattila, D. Interchange of humpback whales in Oceania (South Pacific), 1999 to 2004. Paper SC/A06/HW55 presented to IWC Comprehensive Assessment of Southern Hemisphere Humpback whales 2006, Hobart, Australia.

- Garrigue, C.; Franklin, T.; Russell, K.; Burns, D.; Poole, M.; Paton, D.; Hauser, N.; Oremus, M.; Constantine, R.; Childerhouse, S.; Mattila, D.; Gibbs, N.; Franklin, W.; Robbins, J.; Clapham, P.; Baker, C.S. 2007. First assessment of interchange of humpback whales between Oceania and the east coast of Australia. Paper SC/59/SH15 presented to the Scientific Committee of the International Whaling Commission Annual Meeting 2007, Anchorage, Alaska.
- Gibbs, N.; Paton, D.; Childerhouse, S.; Clapham, P. 2006. Assessment of the current abundance of humpback whales in the Lomaiviti Island Group of Fiji and a comparison with historical data. Paper SC/A06/HW34 presented to IWC Comprehensive Assessment of Southern Hemisphere Humpback whales 2006, Hobart, Australia.
- International Whaling Commission. 2005. Report of the Scientific Committee. Annex H. Report of the Sub-Committee on Other Southern Hemisphere Whale Stocks. *Journal of Cetacean Research and Management (Supplement) 7*: 236.
- International Whaling Commission. 2007. Report of the Workshop on the Comprehensive Assessment of Southern Hemisphere Humpback Whales. Report SC/58/Rep5 to the International Whaling Commission.
- Jackson, J.; Zerbini, A.; Clapham, P.; Garrigue, C.; Hauser, N.; Poole, M.; Baker, C.S. 2006. A Bayesian assessment of humpback whales on breeding grounds of eastern Australia and Oceania (IWC Stocks, E1, E2, E3 and F). Paper SC/A06/HW52 presented to IWC Comprehensive Assessment of Southern Hemisphere Humpback whales 2006, Hobart, Australia.
- Jackson, J. 2008. Comprehensive Assessment and modeling. Page 10 *in*: Report of the Annual Meeting of the South Pacific Whale Research Consortium. South Pacific Whale Research Consortium, P.O. Box 3069, Avarua, Rarotonga, Cook Islands.
- Mackintosh, N.A. 1942. The southern stocks of whalebone whales. *Disc. Rep.* 22:197-300.
- Mackintosh, N.A. 1965. The stocks of whales. Fishing News (Books) Ltd, London.
- Mikhalev, Y.A. 2000. Biological characteristics of humpbacks taken in Antarctic Area V by the whaling fleets *Slava* and *Sovietskaya Ukraina*. unpublished report to the Scientific Committee of the International Whaling Commission, unpublished SC/52/IA.
- Noad, M.; Cato, D.H.; Paton, D. 2006. Absolute and relative abundance estimates of Australian east coast humpback whales. Paper SC/A06/HW27 presented to IWC Comprehensive Assessment of Southern Hemisphere Humpback whales 2006, Hobart, Australia
- Olavarría, C.; Anderson, M.; Paton, D.; Burns, D.; Brasseur, M.; Garrigue, C.; Hauser, N.; Poole, M.; Caballero, S.; Flórez-González, L.; Baker, C.S. 2006. Eastern Australia humpback whale genetic diversity and their relationship with Breeding Stocks D, E, F and G. Paper SC/58/SH25 presented to the Scientific Committee of the International Whaling Commission Annual Meeting 2006, St Kitts.
- Olavarría, C.; Baker, C.S.; Garrigue, C.; Poole, M.; Hauser, N.; Caballero, S.; Flórez-González, L.; Brasseur, M.; Bannister, J.; Capella, J.; Clapham, P.; Dodemont, R.; Donoghue, M.; Jenner, C.; Jenner, M.; Moro, D.; Oremus, M.; Paton, D.; Russell, K. 2007. Population structure of humpback whales throughout the South Pacific and the origins of the eastern Polynesian breeding grounds. *Marine Ecology - Progress Series* 330:257-268.
- Oosterman, A.; Whicker, M. 2008. Norfolk Island Whale Surveys. Reviewing the observations of 2003-2007. Unpublished Report to South Pacific Whale Research Consortium. South Pacific Whale Research Consortium, P.O. Box 3069, Avarua, Rarotonga, Cook Islands. 9 p.
- Paton, D.; Oosterman, A.; Whicker, M.; Kenny, I. 2006. Preliminary assessment of sighting survey data of humpback whales, Norfolk Island, Australia. Paper SC/A06/HW36 presented to IWC Comprehensive Assessment of Southern Hemisphere Humpback whales 2006, Hobart, Australia.
- Rasmussen, K.; Palacios, D.; Calambokidis, J.; Saborío, M.; Dalla Rosa, L.; Secchi, E.; Steiger, S.; Allen, J.; Stone, G. 2007. Southern Hemisphere humpback whales wintering off Central America: insights from water temperature into the longest mammalian migration. *Biology Letters* 3:302-305.
- Rice, D.W. 1998. *Marine Mammals of the World. Systematics and Distribution*. Special Publication Number 4. The Society for Marine Mammalogy. 231 p.
- South Pacific Whale Research Consortium. 2008. Report of the Annual Meeting of the South Pacific Whale Research Consortium, Auckland, 5-8 February 2008. South Pacific Whale Research Consortium, P.O. Box 3069, Avarua, Rarotonga, Cook Islands. 36 p.
- Taylor, B.L., S.J. Chivers, J. Larese, and W.F. Perrin. 2007. Generation length and percent mature

- estimates for IUCN assessments of cetaceans. Administrative Report LJ-07-01 available from Southwest Fisheries Science Center, National Marine Fisheries Service, 8604 La Jolla Shores Dr., La Jolla, CA 92038 USA.
- Townsend, C.H. 1935. The distribution of certain whales as shown by logbook records of American whaleships. *Zoologica* 19:1-50.
- Waples, R. S. & Gaggiotti, O. 2006. What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology* 15(6): 1419-1439
- Wright, S.1978. *Evolution and the Genetics of Populations; Vol 4. Variability within and among natural populations.* University of Chicago Press.

Appendix 1 International Whaling Commission recognised stock structure of humpback whales in the Southern Hemisphere. Reprinted from IWC 2005.

