

**LIST OF BIRD SPECIES THAT BELONG TO THE FAMILIES AND GENERA CURRENTLY
AGGREGATED UNDER APPENDIX II, MEET THE CMS CRITERIA AND HAVE AN
UNFAVOURABLE CONSERVATION STATUS**

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Anatidae	Emperor Goose	<i>Anser canagicus</i>	A member of a disaggregated family from Appendix II. (Anatidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Anatidae	Long-tailed Duck	<i>Clangula hyemalis</i>	A member of a disaggregated family from CMS Appendix II. (Anatidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Anatidae	Spectacled Eider	<i>Somateria fischeri</i>	A member of a disaggregated family from Appendix II. (Anatidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Anatidae	Common Eider	<i>Somateria mollissima</i>	A member of a disaggregated family from CMS Appendix II. (Anatidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: substantial post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Anatidae	Velvet Scoter	<i>Melanitta fusca</i>	A member of a disaggregated family from CMS Appendix II. (Anatidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Anatidae	Black Scoter	<i>Melanitta americana</i>	A member of a disaggregated family from Appendix II. (Anatidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Anatidae	Scaly-sided Merganser	<i>Mergus squamatus</i>	A member of a disaggregated family from Appendix II. (Anatidae). IUCN Red List Status: Endangered; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km north-south. Meets CMS movement criteria.	EN
Anatidae	Orinoco Goose	<i>Neochen jubata</i>	A member of a disaggregated family from Appendix II. (Anatidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km but in no consistent direction. Meets CMS movement criteria.	NT
Anatidae	White-winged Duck	<i>Asarcornis scutulata</i>	A member of a disaggregated family from Appendix II. (Anatidae). IUCN Red List Status: Endangered; Movements characterised as: Sedentary/resident: breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel <100 km but in no consistent direction. Meets CMS movement criteria.	EN

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Anatidae	Common Pochard	<i>Aythya ferina</i>	A member of a disaggregated family from CMS Appendix II. (Anatidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Vulnerable; Movements characterised as: Partial migrant: substantial post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Anatidae	Spectacled Duck	<i>Specularias specularis</i>	A member of a disaggregated family from Appendix II. (Anatidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: substantial post-breeding range expansion with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km north-south. Meets CMS movement criteria.	NT
Anatidae	Falcated Duck	<i>Mareca falcata</i>	A member of a disaggregated family from Appendix II. (Anatidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Phoenicopteridae	Chilean Flamingo	<i>Phoenicopterus chilensis</i>	A member of a disaggregated family from Appendix II. (Phoenicopteridae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift and moderately nomadic in both breeding and non-breeding seasons; individuals commonly travel >1,000 km north-south; at least some movements primarily altitudinal. Meets CMS movement criteria.	NT
Phoenicopteridae	Lesser Flamingo	<i>Phoeniconaias minor</i>	A member of a disaggregated family from CMS Appendix II. (Phoenicopteridae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Sedentary/resident but strongly nomadic in both breeding and non-breeding seasons; individuals commonly travel >1,000 km but in no consistent direction. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Gruidae	Sarus Crane	<i>Antigone antigone</i>	A member of an aggregated Genus Antigone. IUCN Red List Status: Vulnerable; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km but in no consistent direction; Australian subpopulation does not migrate across international boundaries. Meets CMS movement criteria.	VU
Gruidae	Whooping Crane	<i>Grus americana</i>	A member of a disaggregated genus Grus. IUCN Red List Status: Endangered; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	EN
Pluvianellidae	Magellanic Plover	<i>Pluvianellus socialis</i>	A member of a disaggregated family from Appendix II. (Pluvianellidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: substantial post-breeding range expansion with breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel 100-1,000 km north-south. Meets CMS movement criteria.	NT
Haematopodidae	Eurasian Oystercatcher	<i>Haematopus ostralegus</i>	A member of a disaggregated family from Appendix II. (Haematopodidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Charadriidae	Diademed Plover	<i>Phegornis mitchellii</i>	A member of a disaggregated family from Appendix II. (Charadriidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel 100-1,000 km north-south; at least some movements primarily altitudinal. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Charadriidae	Piping Plover	<i>Charadrius melodus</i>	A member of a disaggregated family from Appendix II. (Charadriidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Charadriidae	White-faced Plover	<i>Charadrius dealbatus</i>	A member of a disaggregated family from Appendix II. (Charadriidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel 100-1,000 km north-south. Meets CMS movement criteria.	DD
Charadriidae	Snowy Plover	<i>Charadrius nivosus</i>	A member of a disaggregated family from Appendix II. (Charadriidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: substantial post-breeding range shift: breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel >1,000 km north-south; at least one subpopulation separate from the migration pathway considered sedentary. Meets CMS movement criteria.	NT
Charadriidae	Chestnut-banded Plover	<i>Charadrius pallidus</i>	A member of a disaggregated family from CMS Appendix II. (Charadriidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift and moderately nomadic in both breeding and non-breeding seasons; individuals commonly travel 100-1,000 km north-south. Meets CMS movement criteria.	NT
Charadriidae	Mountain Plover	<i>Charadrius montanus</i>	A member of a disaggregated family from Appendix II. (Charadriidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: substantial post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south; at least one subpopulation separate from the migration pathway considered sedentary. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Charadriidae	Northern Lapwing	<i>Vanellus vanellus</i>	A member of a disaggregated family from CMS Appendix II. (Charadriidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: substantial post-breeding range shift: breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Scolopacidae	Eurasian Curlew	<i>Numenius arquata</i>	A member of a disaggregated family from CMS Appendix II. (Scolopacidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Scolopacidae	Bar-tailed Godwit	<i>Limosa lapponica</i>	A member of a disaggregated family from CMS Appendix II. (Scolopacidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Scolopacidae	Black-tailed Godwit	<i>Limosa limosa</i>	A member of a disaggregated family from CMS Appendix II. (Scolopacidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Scolopacidae	Curlew Sandpiper	<i>Calidris ferruginea</i>	A member of a disaggregated family from CMS Appendix II. (Scolopacidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Scolopacidae	Red-necked Stint	<i>Calidris ruficollis</i>	A member of a disaggregated family from Appendix II. (Scolopacidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Scolopacidae	Asian Dowitcher	<i>Limnodromus semipalmatus</i>	A member of a disaggregated family from Appendix II. (Scolopacidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Scolopacidae	Fuegian Snipe	<i>Gallinago stricklandii</i>	A member of a disaggregated family from Appendix II. (Scolopacidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel 100-1,000 km north-south; at least one subpopulation separate from the migration pathway considered sedentary. Meets CMS movement criteria.	NT
Scolopacidae	Wood Snipe	<i>Gallinago nemoricola</i>	A member of a disaggregated family from Appendix II. (Scolopacidae). IUCN Red List Status: Vulnerable; Movements characterised as: Partial migrant: post-breeding range shift with breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel 100-1,000 km north-south; at least some movements primarily altitudinal. Meets CMS movement criteria.	VU
Scolopacidae	Great Snipe	<i>Gallinago media</i>	A member of a disaggregated family from CMS Appendix II. (Scolopacidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Scolopacidae	Grey-tailed Tattler	<i>Tringa brevipes</i>	A member of a disaggregated family from Appendix II. (Scolopacidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Accipitridae	Bearded Vulture	<i>Gypaetus barbatus</i>	A member of a disaggregated family from CMS Appendix II. (Accipitridae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Sedentary/resident: both breeding and non-breeding locations reliably occupied every season; immatures may migrate beyond normal adult range before returning to breed. Meets CMS movement criteria.	NT
Accipitridae	Bateleur	<i>Terathopius ecaudatus</i>	A member of a disaggregated family from Appendix II. (Accipitridae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km north-south; immatures may migrate beyond normal adult range before returning to breed; at least one subpopulation separate from the migration pathway considered sedentary. Meets CMS movement criteria.	NT
Accipitridae	Beaudouin's Snake-eagle	<i>Circaetus beaudouini</i>	A member of a disaggregated family from Appendix II. (Accipitridae). IUCN Red List Status: Vulnerable; Movements characterised as: Partial migrant: substantial post-breeding range shift: breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel 100-1,000 km north-south. Meets CMS movement criteria.	VU
Accipitridae	Southern Banded Snake-eagle	<i>Circaetus fasciolatus</i>	A member of a disaggregated family from Appendix II. (Accipitridae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range expansion with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km north-south. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Accipitridae	Himalayan Griffon	<i>Gyps himalayensis</i>	A member of a disaggregated family from CMS Appendix II. (Accipitridae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel <100 km but in no consistent direction; at least some movements primarily altitudinal; immatures may migrate beyond normal adult range before returning to breed. Meets CMS movement criteria.	NT
Accipitridae	Cinereous Vulture	<i>Aegypius monachus</i>	A member of a disaggregated family from CMS Appendix II. (Accipitridae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift and substantial expansion and moderately nomadic in both breeding and non-breeding seasons; individuals commonly travel >1,000 km north-south; at least one subpopulation separate from the migration pathway considered sedentary. Meets CMS movement criteria.	NT
Accipitridae	Tawny Eagle	<i>Aquila rapax</i>	A member of a disaggregated family from CMS Appendix II. (Accipitridae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Vulnerable; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km north-south; at least one subpopulation separate from the migration pathway considered sedentary. Meets CMS movement criteria.	VU

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Accipitridae	Black Harrier	<i>Circus maurus</i>	A member of a disaggregated family from CMS Appendix II. (Accipitridae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Endangered; Movements characterised as: Partial migrant: substantial post-breeding range shift: breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel >1,000 km north-south; sometimes irrupts beyond normal range in significant numbers; at least one subpopulation separate from the migration pathway considered sedentary. Meets CMS movement criteria.	EN
Accipitridae	Pallid Harrier	<i>Circus macrourus</i>	A member of a disaggregated family from CMS Appendix II. (Accipitridae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Accipitridae	Grey-bellied Goshawk	<i>Accipiter poliogaster</i>	A member of a disaggregated family from Appendix II. (Accipitridae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Accipitridae	Lesser Fish-eagle	<i>Icthyophaga humilis</i>	A member of a disaggregated family from Appendix II. (Accipitridae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km but in no consistent direction; at least some movements primarily altitudinal; at least one subpopulation separate from the migration pathway considered sedentary. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Accipitridae	Red Kite	<i>Milvus milvus</i>	A member of a disaggregated family from CMS Appendix II. (Accipitridae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km north-south. Meets CMS movement criteria.	NT
Falconidae	Striated Caracara	<i>Phalcoboenus australis</i>	A member of a disaggregated family from Appendix II. (Falconidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km north-south; at least some movements primarily altitudinal. Meets CMS movement criteria.	NT
Falconidae	Red-headed Falcon	<i>Falco chicquera</i>	A member of a disaggregated family from Appendix II. (Falconidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift and moderately nomadic in both breeding and non-breeding seasons; individuals commonly travel 100-1,000 km but in no consistent direction; at least one subpopulation separate from the migration pathway considered sedentary. Meets CMS movement criteria.	NT
Falconidae	Sooty Falcon	<i>Falco concolor</i>	A member of a disaggregated family from CMS Appendix II. (Falconidae). Although currently listed on Appendix II database as a species, not officially inscribed. IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Oriolidae	Silver Oriole	<i>Oriolus mellianus</i>	A member of a disaggregated family from Appendix II. (Oriolidae, formerly Muscicapidae). IUCN Red List Status: Endangered; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	EN

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Vireonidae	Black-capped Vireo	<i>Vireo atricapilla</i>	A member of a disaggregated family from Appendix II. (Vireonidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Monarchidae	Japanese Paradise-flycatcher	<i>Terpsiphone atrocaudata</i>	A member of a disaggregated family from Appendix II. (Monarchidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: substantial post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Chaetopidae	Drakensberg Rockjumper	<i>Chaetops aurantius</i>	A member of a disaggregated family from Appendix II. (Chaetopidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range expansion with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel <100 km but in no consistent direction; at least some movements primarily altitudinal. Meets CMS movement criteria.	NT
Acrocephalidae	Large-billed Reed-warbler	<i>Acrocephalus orinus</i>	A member of a disaggregated family from Appendix II. (Acrocephalidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	DD
Acrocephalidae	White-browed Reed-warbler	<i>Acrocephalus tangorum</i>	A member of a disaggregated family from Appendix II. (Acrocephalidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Locustellidae	Marsh Grassbird	<i>Locustella pryeri</i>	A member of a disaggregated family from Appendix II. (Locustellidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Locustellidae	Pleske's Grasshopper-warbler	<i>Locustella pleskei</i>	A member of a disaggregated family from Appendix II. (Locustellidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Locustellidae	Long-billed Grasshopper-warbler	<i>Locustella major</i>	A member of a disaggregated family from Appendix II. (Locustellidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel <100 km but in no consistent direction; at least some movements primarily altitudinal. Meets CMS movement criteria.	NT
Locustellidae	Bristled Grassbird	<i>Chaetornis striata</i>	A member of a disaggregated family from Appendix II. (Locustellidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Partial migrant: post-breeding range shift and strongly nomadic in both breeding and non-breeding seasons; individuals commonly travel 100-1,000 km north-south. Meets CMS movement criteria.	VU
Phylloscopidae	Tytler's Leaf-warbler	<i>Phylloscopus tytleri</i>	A member of a disaggregated family from Appendix II. (Phylloscopidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Phylloscopidae	Ijima's Leaf-warbler	<i>Phylloscopus ijimae</i>	A member of a disaggregated family from Appendix II. (Phylloscopidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Sylviidae	Bush Blackcap	<i>Sylvia nigricapillus</i>	A member of a disaggregated family from Appendix II. (Sylviidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Partial migrant: post breeding range contraction with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km but in no consistent direction; at least some movements primarily altitudinal. Meets CMS movement criteria.	VU
Sylviidae	Dartford Warbler	<i>Sylvia undata</i>	A member of a disaggregated family from Appendix II. (Sylviidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift and substantial expansion with breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel 100-1,000 km but in no consistent direction; at least some movements primarily altitudinal. Meets CMS movement criteria.	NT
Pellorneidae	Rufous-vented Grass-babbler	<i>Laticilla burnesii</i>	A member of a disaggregated family from Appendix II. (Pellorneidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range expansion with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel <100 km but in no consistent direction. Meets CMS movement criteria.	NT
Turdidae	Wood Thrush	<i>Hylocichla mustelina</i>	A member of a disaggregated family from Appendix II. (Turdidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Turdidae	Bicknell's Thrush	<i>Catharus bicknelli</i>	A member of a disaggregated family from Appendix II. (Turdidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Turdidae	Redwing	<i>Turdus iliacus</i>	A member of a disaggregated family from Appendix II. (Turdidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: substantial post-breeding range shift: breeding locations reliably occupied every season but strongly nomadic in non-breeding season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Turdidae	Grey-sided Thrush	<i>Turdus feae</i>	A member of a disaggregated family from Appendix II. (Turdidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Muscicapidae	Zappey's Flycatcher	<i>Cyanoptila cumatilis</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Muscicapidae	Brown-chested Jungle-flycatcher	<i>Cyornis brunneatus</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Muscicapidae	Rusty-bellied Shortwing	<i>Brachypteryx hyperythra</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel <100 km but in no consistent direction; at least some movements primarily altitudinal. Meets CMS movement criteria.	NT
Muscicapidae	Rufous-headed Robin	<i>Larivora ruficeps</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Endangered; Movements characterised as: Full migrant: breeding locations reliably occupied every season but no information on non-breeding movements; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	EN
Muscicapidae	Firethroat	<i>Calliope pectardens</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Muscicapidae	Blackthroat	<i>Calliope obscura</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: breeding locations reliably occupied every season but no information on non-breeding movements; individuals commonly travel >1,000 km north-south; at least some movements primarily altitudinal. Meets CMS movement criteria.	VU
Muscicapidae	Kashmir Flycatcher	<i>Ficedula subrubra</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Muscicapidae	Sentinel Rock-thrush	<i>Monticola explorator</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: post-breeding range expansion with breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel <100 km but in no consistent direction; at least some movements primarily altitudinal. Meets CMS movement criteria.	NT
Muscicapidae	White-browed Bushchat	<i>Saxicola macrorhynchus</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant and moderately nomadic in breeding season and strongly so in non-breeding season; individuals commonly travel 100-1,000 km but in no consistent direction. Meets CMS movement criteria.	VU
Muscicapidae	White-throated Bushchat	<i>Saxicola insignis</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Muscicapidae	Sombre Rockchat	<i>Oenanthe dubia</i>	A member of a disaggregated family from Appendix II. (Muscicapidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Sedentary/resident: breeding locations reliably occupied every season but moderately nomadic in non-breeding season; individuals commonly travel 100-1,000 km but in no consistent direction. Meets CMS movement criteria.	DD
Motacillidae	Meadow Pipit	<i>Anthus pratensis</i>	A member of a disaggregated family from Appendix II. (Motacillidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Partial migrant: substantial post-breeding range shift with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south; at least some movements primarily altitudinal. Meets CMS movement criteria.	NT

Family	Common name	Scientific name	Notes for Appendix II (Sheets 1 or 2)	2018 IUCN Red List Category
Motacillidae	Sprague's Pipit	<i>Anthus spragueii</i>	A member of a disaggregated family from Appendix II. (Motacillidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	VU
Motacillidae	Mountain Pipit	<i>Anthus hoeschi</i>	A member of a disaggregated family from Appendix II. (Motacillidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel >1,000 km north-south. Meets CMS movement criteria.	NT
Motacillidae	Yellow-breasted Pipit	<i>Hemimacronyx chloris</i>	A member of a disaggregated family from Appendix II. (Motacillidae, formerly Muscicapidae). IUCN Red List Status: Vulnerable; Movements characterised as: Partial migrant: substantial post-breeding range expansion with both breeding and non-breeding locations reliably occupied every season; individuals commonly travel 100-1,000 km east-west; at least some movements primarily altitudinal. Meets CMS movement criteria.	VU
Motacillidae	Mekong Wagtail	<i>Motacilla samveasnae</i>	A member of a disaggregated family from Appendix II. (Motacillidae, formerly Muscicapidae). IUCN Red List Status: Near Threatened; Movements characterised as: Full migrant: both breeding and non-breeding locations reliably occupied every season; individuals commonly travel <100 km but in no consistent direction. Meets CMS movement criteria.	NT